Welcome to STN International! Enter x:x

LOGINID:SSSPTA1636DXS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * * * * * * Welcome to STN International * * * * * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America

NEWS 2 Apr 08 "Ask CAS" for self-help around the clock

NEWS 3 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area

NEWS 4 Apr 09 ZDB will be removed from STN

NEWS 5 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB

NEWS 6 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS

NEWS 7 Apr 22 BIOSIS Gene Names now available in TOXCENTER

NEWS 8 Apr 22 Federal Research in Progress (FEDRIP) now available

NEWS 9 Jun 03 New e-mail delivery for search results now available

NEWS 10 Jun 10 MEDLINE Reload

NEWS 11 Jun 10 PCTFULL has been reloaded

NEWS 12 Jul 02 FOREGE no longer contains STANDARDS file segment

NEWS 13 Jul 22 USAN to be reloaded July 28, 2002; saved answer sets no longer valid

NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY

NEWS 15 Jul 30 NETFIRST to be removed from STN

NEWS 16 Aug 08 CANCERLIT reload

NEWS 17 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN

NEWS 18 Aug 08 NTIS has been reloaded and enhanced

NEWS 19 Aug 09 JAPIO to be reloaded August 25, 2002

NEWS 20 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN

NEWS 21 Aug 19 IFIPAT, IFICDB, and IFIUDB have been reloaded NEWS 22 Aug 19 The MEDLINE file segment of TOXCENTER has been reloaded

NEWS 23 Aug 26 Sequence searching in REGISTRY enhanced

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,

CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),

AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS INTER General Internet Information

NEWS LOGIN Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access to STN

NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002

=> file medline biosis caplus

COST IN U.S. DOLLARS SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST 0.21 0.21

FILE 'MEDLINE' ENTERED AT 10:28:13 ON 01 SEP 2002

FILE 'BIOSIS' ENTERED AT 10:28:13 ON 01 SEP 2002 COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s dry eye or xerosis or hypolacrimia or keratoconjuctivitis sicca L1 4177 DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SICCA

=> s animal model

L2 251329 ANIMAL MODEL

=> s 11 and 12

L3 49 L1 AND L2

=> dup rem 13

PROCESSING COMPLETED FOR L3

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

=> d ti so 1-43

L4 ANSWER 1 OF 43 MEDLINE DUPLICATE 1
TI Androgen deficiency, Meibomian gland dysfunction, and

evaporative

dry eye.

SO ANNALS OF THE NEW YORK ACADEMY OF SCIENCES, (2002 Jun) 966 211-22. Ref:

121

Journal code: 7506858. ISSN: 0077-8923.

L4 ANSWER 2 OF 43 MEDLINE

DUPLICATE 2

TI Expression of interleukin-4 in the epidermis of transgenic mice results in

a pruritic inflammatory skin disease: an experimental animal model to study atopic dermatitis.

SO JOURNAL OF INVESTIGATIVE DERMATOLOGY, (2001 Oct) 117 (4) 977-83.

Journal code: 0426720. ISSN: 0022-202X.

L4 ANSWER 3 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Estrogen and progesterone influence on gene expression in the lacrimal

gland.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S712. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

L4 ANSWER 4 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Effect of UTP and cAMP on the fluid secretion in adenovirus type 5 infected pigmented rabbit conjunctiva.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S485. print.

Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04,

L4 ANSWER 5 OF 43 MEDLINE

DUPLICATE 3

TI Sjogren's syndrome: immunological response underlying the disease.

SO ARCHIVUM IMMUNOLOGIAE ET THERAPIAE EXPERIMENTALIS, (2001) 49 (5) 353-60.

Ref: 62

Journal code: 0114365. ISSN: 0004-069X.

- L4 ANSWER 6 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Epithelial mucins of the ocular surface: Structure, biosynthesis and function.
- SO Experimental Eye Research, (September, 2001) Vol. 73, No. 3, pp. 281-289.

print.

ISSN: 0014-4835.

- L4 ANSWER 7 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI A new dry-eye animal model using GVH reactions.
- SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S263. print. Meeting Info.: Annual Meeting of the Association for Research in Vision
- and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001
- L4 ANSWER 8 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Long-term changes in the ocular surface and tear flow after pre-ganglionic, parasympathetic denervation in the rabbit.
- SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S263. print.
- Meeting Info.: Annual Meeting of the Association for Research in Vision
- and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001
- L4 ANSWER 9 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI A new animal model of dry eyes.
- SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S262. print.

 Meeting Info.: Annual Meeting of the Association for Research in Vision
- and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001
- L4 ANSWER 10 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Over-expression of nitric oxide by pro-inflammatory cytokines in cultured

lacrimal gland acinar cells: A possible factor in dry eye syndrome.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S258. print.
Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

- L4 ANSWER 11 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Corneal protection by the ocular mucin secretagogue 15(S)-HETE in a rabbit

dry eye model.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S177. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

- L4 ANSWER 12 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Improvement of corneal barrier function by the P2Y2 agonist

INS365 in a

rat dry eye model.

SO IOVS, (January, 2001) Vol. 42, No. 1, pp. 96-100. print.

- L4 ANSWER 13 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Effects of OPC-12759 ophthalmic suspension on N-acetylcysteine-treated

eyes.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S37. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

- L4 ANSWER 14 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Experimentally induced dry eye produces ocular surface inflammation and epithelial disease.
- SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S31. print.

 Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

- L4 ANSWER 15 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Proliferation of lacrimal gland acinar cells in primary culture. Stimulation by extracellular matrix, EGF, and DHT.
- SO Experimental Eye Research, (May, 2000) Vol. 70, No. 5, pp. 639-649. print.

ISSN: 0014-4835.

- L4 ANSWER 16 OF 43 MEDLINE
- TI Microvascular submandibular gland transfer for severe cases of keratoconjunctivitis sicca.
- SO PLASTIC AND RECONSTRUCTIVE SURGERY, (2000 Sep) 106 (3) 554-60; discussion

561-2.

Journal code: 1306050. ISSN: 0032-1052.

- L4 ANSWER 17 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Gefarnate stimulates goblet cell repopulation after an experimental wound

to the tarsal conjunctiva in dry eye rabbits.

SO IOVS, (March 15, 2000) Vol. 41, No. 4, pp. S275. print.. Meeting Info.: Annual Meeting of the Association in Vision and Opthalmology. Fort Lauderlade, Florida, USA April 30-May 05, 2000

Association for Research in Vision and Ophthalmology

- L4 ANSWER 18 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Comparative study of the efficacy of various moisturizers on the skin of

the ASR miniature swine.

SO Skin Pharmacology and Applied Skin Physiology, (March April, 2000) Vol.

13, No. 2, pp. 120-127.

ISSN: 1422-2868.

- L4 ANSWER 19 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Pregnancy and lactation induce changes in FGF2 and TGFbeta1 distribution

in the rabbit lacrimal gland: A potential basis for predisposition to dry eye syndrome.

SO Journal of Investigative Medicine, (January, 2000) Vol. 48, No. 1, pp.

58A. print.

Meeting Info.: Meeting of the American Federation for Medical Research,

Western Region Carmel, California, USA February 09-12, 2000 ISSN: 1081-5589.

L4 ANSWER 20 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Autologous lacrimal-lymphoid mixed-cell reactions induce dacryoadenitis in

rabbits.

SO Experimental Eye Research, (July, 2000) Vol. 71, No. 1, pp. 23-31. print.

ISSN: 0014-4835.

L4 ANSWER 21 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Changes in expression of Maxi-K channels in aging NZB/NZW F1 female mouse

lacrimal gland acinar cells.

SO IOVS, (March 15, 1999) Vol. 40, No. 4, pp. S985.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA May 9-14, 1999 Association

for Research in Vision and Opthalmology

•

L4 ANSWER 22 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Functional recovery and prevention of apoptosis in lacrimal gland acinar

cells by topical application of cyclosporine A in Sjogren's syndrome model

mouse.

SO IOVS, (March 15, 1999) Vol. 40, No. 4, pp. S554.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA May 9-14, 1999 Association

for Research in Vision and Opthalmology

.

L4 ANSWER 23 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Removal of main and accessory lacrimal glands in experimental dry eye models in rabbits and squirrel monkeys.

SO IOVS, (March 15, 1999) Vol. 40, No. 4, pp. S538.

Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA May 9-14, 1999 Association

for Research in Vision and Opthalmology

٠

L4 ANSWER 24 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Impact of gender on exocrine gland inflammation in mouse models of

Sjogren's syndrome.

SO Experimental Eye Research, (Oct., 1999) Vol. 69, No. 4, pp. 355-366.

ISSN: 0014-4835.

L4 ANSWER 25 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Development of a simple **dry** eye model in the albino rabbit and evaluation of some tear substitutes.

SO Ophthalmic Research, (May-June, 1999) Vol. 31, No. 3, pp. 229-235.

ISSN: 0030-3747.

L4 ANSWER 26 OF 43 CAPLUS COPYRIGHT 2002 ACS

TI Transfer of human serum IgG to nonobese diabetic Ig.mu.null mice reveals a

role for autoantibodies in the loss of secretory function of exocrine tissues in Sjogren's syndrome

SO Proceedings of the National Academy of Sciences of the United States of

America (1998), 95(13), 7538-7543 CODEN: PNASA6; ISSN: 0027-8424

L4 ANSWER 27 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Biochemical analysis of ocular surface mucin abnormalities in dry eye: The canine model.

SO Experimental Eye Research, (Dec., 1998) Vol. 67, No. 6, pp. 709-718

ISSN: 0014-4835.

L4 ANSWER 28 OF 43 CAPLUS COPYRIGHT 2002 ACS

TI The immunology of Sjoegren's syndrome: emphasis on the role of salivary

gland autoantigen

SO Biomedical Reviews (1998), 9, 131-141 CODEN: BMREES; ISSN: 1310-392X

L4 ANSWER 29 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Lacrimal gland innervation is not altered with the onset and progression

of disease in a murine model of Sjogren's syndrome.

SO Clinical Immunology and Immunopathology, (Nov., 1998) Vol. 89, No. 2, pp.

126-133.

ISSN: 0090-1229.

L4 ANSWER 30 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term dry eye model.

SO Journal of Ocular Pharmacology and Therapeutics, (April, 1998) Vol. 14,

No. 2, pp. 99-107.

ISSN: 1080-7683.

L4 ANSWER 31 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Gefarnate stimulates secretion of mucin-like glycoproteins by corneal

epithelium in vitro and protects corneal epithelium from desiccation in

vivo.

SO Experimental Eye Research, (1997) Vol. 65, No. 4, pp. 569-574. ISSN: 0014-4835.

L4 ANSWER 32 OF 43 MEDLINE

TI Sjogren's syndrome. Controversies and progress.

SO CLINICS IN LABORATORY MEDICINE, (1997 Sep) 17 (3) 431-44. Ref: 78

Journal code: 8100174. ISSN: 0272-2712.

L4 ANSWER 33 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The effects of artificial tear solutions on wound healing in full thickness corneal incisions.

SO Acta Physiologica Hungarica, (1997) Vol. 85, No. 3, pp. 251-258. ISSN: 0231-424X.

L4 ANSWER 34 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Androgen stimulation of lacrimal gland function in mouse models of Siogren's syndrome.

SO Journal of Steroid Biochemistry and Molecular Biology, (1997) Vol. 60, No.

3-4, pp. 237-245.

ISSN: 0960-0760.

L4 ANSWER 35 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Non-Sjogren dry eye: Pathogenesis diagnosis and

animal models.

SO Sullivan, D. A. [Editor]. Advances in Experimental Medicine and

(1994) Vol. 350, pp. 471-488. Advances in Experimental Medicine

Biology; Lacrimal gland, tear film, and dry eye syndromes: Basic science

and clinical relevance.

Publisher: Plenum Press 233 Spring Street, New York, New York,

Meeting Info.: International Conference Southampton, Bermuda November

14-17, 1992

ISSN: 0065-2598. ISBN: 0-306-44676-6.

L4 ANSWER 36 OF 43 MEDLINE

TI Epidemiology, pathogenesis, animal models, and treatment of Sjogren's syndrome.

SO CURRENT OPINION IN RHEUMATOLOGY, (1994 Sep) 6 (5) 501-8. Ref: 89

Journal code: 9000851. ISSN: 1040-8711.

L4 ANSWER 37 OF 43 MEDLINE

TI Non-Sjogren dry eye: pathogenesis diagnosis and animal models.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1994) 350 471-88. Ref:

127

Journal code: 0121103. ISSN: 0065-2598.

L4 ANSWER 38 OF 43 MEDLINE **DUPLICATE 4**

TI Efficacy of laser punctal occlusion.

SO OPHTHALMOLOGY, (1992 Apr) 99 (4) 618-21.

Journal code: 7802443. ISSN: 0161-6420.

L4 ANSWER 39 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI COMPARATIVE EFFECTS OF PRESERVATIVE FREE TEAR SUBSTITUTES ON THE RABBIT

CORNEA A SCANNING ELECTRON MICROSCOPIC EVALUATION.

SO ANNUAL SPRING MEETING OF THE ASSOCIATION FOR RESEARCH IN VISION AND

OPHTHALMOLOGY, SARASOTA, FLORIDA, USA, APRIL 28-MAY 3, 1991. INVEST

OPHTHALMOL VISUAL SCI. (1991) 32 (4), 733.

CODEN: IOVSDA. ISSN: 0146-0404.

L4 ANSWER 40 OF 43 MEDLINE

TI Argon laser versus thermal cautery for punctal occlusion. An animal

SO OPHTHALMIC PLASTIC AND RECONSTRUCTIVE SURGERY, (1991) 7 (3) 173-6.

Journal code: 8508431. ISSN: 0740-9303.

L4 ANSWER 41 OF 43 MEDLINE

TI Ocular disease associated with Epstein-Barr virus infection.

SO SURVEY OF OPHTHALMOLOGY, (1990 Sep-Oct) 35 (2) 145-

50. Ref: 59

Journal code: 0404551. ISSN: 0039-6257.

L4 ANSWER 42 OF 43 MEDLINE **DUPLICATE 5**

TI Sjogren's syndrome in MRL/1 and MRL/n mice.

SO ARTHRITIS AND RHEUMATISM, (1984 Feb) 27 (2) 157-65. Journal code: 0370605. ISSN: 0004-3591.

L4 ANSWER 43 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI A NEW MOUSE CORNEAL DYSTROPHY.

SO ANNUAL MEETING OF THE ASSOCIATION FOR RESEARCH IN VISION AND OPHTHALMOLOGY

INCORPORATED, SARASOTA, FLA., USA, APR. 30-MAY 4, 1979. INVEST OPHTHALMOL

VISUAL SCI. (1979) 0 (SUPPL), 74.

CODEN: IOVSDA. ISSN: 0146-0404.

=> d ibib ab 36,35,30,27,25,24,17,14,1211,9,7 43 ANSWERS ARE AVAILABLE. SPECIFIED ANSWER NUMBER EXCEEDS ANSWER SET SIZE The answer numbers requested are not in the answer set. ENTER ANSWER NUMBER OR RANGE (1):end

=> d ibib ab 36,35,30,27,25,24,17,14,12,11,9,7

L4 ANSWER 36 OF 43 MEDLINE

ACCESSION NUMBER: 95085859 MEDLINE

DOCUMENT NUMBER: 95085859 PubMed ID: 7993708 Epidemiology, pathogenesis, animal models , and treatment of Sjogren's syndrome.

AUTHOR: Fox R I

CORPORATE SOURCE: Department of Rheumatology, Scripps Clinic and Research

Foundation, La Jolla, CA 92037.

CONTRACT NUMBER: M01 RR00833 (NCRR)

SOURCE: CURRENT OPINION IN RHEUMATOLOGY, (1994 Sep) 6 (5) 501-8.

Ref: 89

Journal code: 9000851. ISSN: 1040-8711.

United States PUB. COUNTRY:

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

General Review; (REVIEW) (REVIEW, TUTORIAL)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 199501

ENTRY DATE: Entered STN: 19950126 Last Updated on STN: 19950126 Entered Medline: 19950119

AB Sjogren's syndrome is characterized by dry eyes

(xerophthalmia) and dry mouth (xerostomia). Although general agreement

exists about the ocular features of Sjogren's syndrome, significant controversy surrounds the classification criteria for defining the oral component. This has resulted in confusion in both the clinical and the research literature. The recent litigation involving silicone breast implants has forced clinicians to evaluate critically whether the condition of a particular patient fulfills specific diagnostic criteria for Sjogren's syndrome. Research studies have focused on potential genetic

and environmental factors in the pathogenesis of Sjogren's syndrome. Studies of various ethnic populations have demonstrated an association

with particular HLA-DR and -DQ alleles. However, the associated alleles

are different for each ethnic group. No single environmental agent has

been identified as a critical cofactor. Studies have concentrated on herpesviruses (Epstein-Barr virus and human herpesvirus type 6),

C virus, and retroviruses. Epstein-Barr virus isolated from patients

altered the ability to transform and lytically infect particular types of lymphocytes. Hepatitis C can lead to sicca symptoms, even in patients with

relatively normal salivary gland biopsy findings. One report of Japanese

patients indicated the presence of human T cell lymphotropic virus type

I-like tax genes in the salivary biopsy specimens of a subset of

and no gag, pol, or env sequences; this finding suggested a potential infection by a defective retrovirus. Studies on the pathogenesis have indicated that cytokines produced in the salivary gland are similar to

helper type 1 lymphocytes (interferon gamma).(ABSTRACT TRUNCATED AT 250

WORDS)

L4 ANSWER 35 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL

ABSTRACTS INC.

ACCESSION NUMBER: 1995:1295 BIOSIS
DOCUMENT NUMBER: PREV199598015595
TITLE: Non-Sjogren dry eye: Pathogenesis

diagnosis and animal models.

AUTHOR(S): Bron, Anthony J.

CORPORATE SOURCE: Nuffield Lab. Ophthalmol., Univ. Oxford,

Oxford OX2 6AW UK

SOURCE: Sullivan, D. A. [Editor]. Advances in Experimental

Medicine

and Biology, (1994) Vol. 350, pp. 471-488. Advances in Experimental Medicine and Biology; Lacrimal gland, tear film, and dry eye syndromes: Basic science and clinical relevance.

Publisher: Plenum Press 233 Spring Street, New York, New

York, USA.

Meeting Info.: International Conference Southampton,

Bermuda November 14-17, 1992

ISSN: 0065-2598. ISBN: 0-306-44676-6.

DOCUMENT TYPE: Book; Conference

LANGUAGE: English

L4 ANSWER 30 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1998:261667 BIOSIS DOCUMENT NUMBER: PREV199800261667

TITLE: Lactoferrin suppresses loss of corneal epithelial

integrity

in a rabbit short-term dry eye model.

AUTHOR(S): Fujihara, Tsutomu (1); Nagano, Takashi; Nakamura, Masatsugu; Shirasawa, Eiichi

CORPORATE SOURCE: (1) Santen Pharmaceutical Co., Ltd. Nara Res. Develop.

Cent., 8916-16, Takayama-cho, Ikoma-shi, Nara, 630-01

Japan

SOURCE: Journal of Ocular Pharmacology and Therapeutics,

(April,

1998) Vol. 14, No. 2, pp. 99-107.

ISSN: 1080-7683.
DOCUMENT TYPE: Article

LANGUAGE: English
AB Human tear fluid contains lactoferrin at the highest concentration.

patients with **dry** eye such as Sjogren's syndrome, the concentration of lactoferrin in the tears is approximately half the normal

value. The present study utilizes a short-term rabbit dry
eye model to evaluate if lactoferrin containing eye drops can
reverse any of the damage produced by blockage of blinking with an
ocular

speculum. Damage was evaluated based on the extent of methylene blue

staining in histological sections. After 3 h of desiccation, the amount of extractable dye recovered following sacrifice increased by more than

4-fold in the vehicle-treated eyes. However, in those rabbits treated with

1% lactoferrin, dye recovery was only 40% of the value in the vehicle-treated eyes. Between 1-3 h and over a concentration range from

0.01 to 1% lactoferrin, the decreases in staining were both time and concentration dependent. Alternatively, if 1% lactoferrin was applied during the desiccation period, there was partial restoration of corneal epithelial integrity. These results suggest that lactoferrin may be of therapeutic value in decreasing the loss of corneal epithelial integrity in dry eye.

L4 ANSWER 27 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1999:151081 BIOSIS DOCUMENT NUMBER: PREV199900151081

TITLE: Biochemical analysis of ocular surface mucin

abnormalities

in dry eye: The canine model.

AUTHOR(S): Hicks, Sally J.; Corfield, Anthony P.; Kaswan, Renee L.;

Hirsh, Susan; Stern, Michael; Bara, Jacques; Carrington, Stephen D. (1)

CORPORATE SOURCE: (1) Dep. Anat., Sch. Vet. Sci., Southwell St., Bristol BS2

8EJ UK

SOURCE: Experimental Eye Research, (Dec., 1998) Vol. 67,

No. 6, pp.

709-718.

ISSN: 0014-4835.

DOCUMENT TYPE: Article

LANGUAGE: English

AB This study examines the canine model of keratoconjunctivitis sicca (KCS, '

dry eye') in order to establish the biochemical basis of altered ocular mucin secretion in this condition. It follows a previous examination of ocular mucins in the normal dog. Mucus was collected by

suction from the ocular surface of dogs with KCS, and dispersed in guanidine hydrochloride containing a cocktail of protease inhibitors. Caesium chloride density gradient centrifugation was used to separate floating 'rafts' of cell membranes from gradients containing secreted mucins. Gradient fractions were collected into pools on the basis of differential staining by Periodic Acid Schiff, Wheat Germ

Agglutinin, and

antibodies to MUC5AC peptide. High molecular weight glycoproteins were

purified from the pooled material by gel filtration chromatography.

Membrane-associated glycoproteins were also derived from the
membrane

rafts using octyl glucoside extraction and/or reduction and alkylation. Secreted mucins and membrane extracts from KCS samples were compared to

equivalent material obtained from normal eyes. Density gradient staining

profiles for normal and KCS mucus were similar over the buoyant density

range typical for secreted mucins, enabling the collection of identical pools of gradient fractions for direct comparison. The following differences were observed in KCS secreted mucins compared to normal

samples: an increase in the proportion of mucin with low buoyant density;

a decrease in mannose content detected with Concanavalin A lectin; an

increase in N-acetylglucosamine structures detected with Lycopersicon

esculentum lectin; increased migration and lack of evidence for distinct

subunit structure on agarose gets. In membrane extracts, the main difference was the presence of T antigen (Galbeta1-3GalNAc) in KCS. These

results demonstrate alterations in the subunit linkage of mucins in KCS,

and suggest that glycosylation, core protein expression and/or post-synthetic modification of ocular surface mucins may also be changed.

L4 ANSWER 25 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1999:307975 BIOSIS DOCUMENT NUMBER: PREV199900307975

TITLE: Development of a simple **dry** eye model in the albino rabbit and evaluation of some tear substitutes.

AUTHOR(S): Burgalassi, Susi (1); Panichi, Luana; Chetoni, Patrizia:

Saettone, M. Fabrizio; Boldrini, Enrico

CORPORATE SOURCE: (1) Department of Pharmaceutical Sciences, University of

Pisa, Via Bonanno 33, I-56126, Pisa Italy

SOURCE: Ophthalmic Research, (May-June, 1999) Vol. 31, No. inflammation was far worse in lacrimal glands of males, whereas immune 3, pp. 229-235. pathology was ISSN: 0030-3747. far greater in salivary tissues in females; and (3) no gender-related variations were present in the degree of inflammation in lacrimal DOCUMENT TYPE: Article LANGUAGE: **English** of bcl-2+/lpr and bcl-2-/lpr mice or in submandibular tissues of SUMMARY LANGUAGE: English AB The present paper is concerned with the development of a simple MRL/lpr, C3H/lpr, bcl-2+/lpr and bcl-2-/lpr mice. Our findings demonstrate dry eye model in the rabbit, induced by daily repeated instillations of 1.0% atropine sulphate. The evolution of the dry that eye syndrome in the animals was assessed by the Schirmer I test gender-, strain- and tissue-related differences exist in the extent of inflammation in several mouse models of Sjogren's syndrome. and by examination of the cornea after fluorescein staining. The L4 ANSWER 17 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL produced rapidly some typical dry eye symptoms and ABSTRACTS INC. could be satisfactorily used for a preliminary assessment of the ACCESSION NUMBER: 2000:263066 BIOSIS protective activity of some polymeric tear substitutes. These were DOCUMENT NUMBER: PREV200000263066 based Gefarnate stimulates goblet cell repopulation after an on hydroxypropylmethylcellulose, sodium hyaluronate, sodium TITLE: experimental wound to the tarsal conjunctiva in dry polyacrylate eye rabbits. or tamarind gum. The latter polymer showed the best overall results. AUTHOR(S): Kim, O. (1); Toshida, H. (1); Hamano, T.; Nakata, Ferning tests on the formulations were also performed: their validity K.; predictors of the efficacy of tear substitutes is discussed. Nakamura, M.; Nguyen, D. (1); Beuerman, R. (1) CORPORATE SOURCE: (1) LSU Eye Center, New Orleans, LA USA IOVS, (March 15, 2000) Vol. 41, No. 4, pp. S275. L4 ANSWER 24 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL SOURCE: ABSTRACTS INC. print.. Meeting Info.: Annual Meeting of the Association in Vision ACCESSION NUMBER: 1999:528292 BIOSIS DOCUMENT NUMBER: PREV199900528292 and Opthalmology. Fort Lauderlade, Florida, USA April 30-May 05, 2000 Association for Research in Vision and TITLE: Impact of gender on exocrine gland inflammation in Ophthalmology mouse models of Sjogren's syndrome. AUTHOR(S): DOCUMENT TYPE: Conference Toda, Ikuko, Sullivan, Benjamin D., Rocha, LANGUAGE: English Eduardo M.; Da Silveira, Lilia A.; Wickham, L. Alexandra; Sullivan, David SUMMARY LANGUAGE: English L4 ANSWER 14 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL CORPORATE SOURCE: (1) Schepens Eye Research Institute, 20 ABSTRACTS INC. Staniford Street, ACCESSION NUMBER: 2001:287143 BIOSIS Boston, MA, 02114 USA DOCUMENT NUMBER: PREV200100287143 SOURCE: Experimental Eye Research, (Oct., 1999) Vol. 69, No. Experimentally induced dry eye produces TITLE: 4, pp. ocular surface inflammation and epithelial disease. 355-366. AUTHOR(S): Dursun, D. (1); Wang, M. (1); Monroy, D. (1); ISSN: 0014-4835. DOCUMENT TYPE: Article Stern, M. E .: Li, D. Q. (1); Lokeshwar, B. (1); Pflugfelder, S. C. (1) LANGUAGE: English SUMMARY LANGUAGE: English CORPORATE SOURCE: (1) Ocular Surface and Tear Center, Bascom AB Sjogren's syndrome is a complex autoimmune disorder, that occurs Palmer Eye almost Institute, Miami, FL USA SOURCE: IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S31. exclusively in females, induces extensive lymphocyte accumulation print. in Meeting Info.: Annual Meeting of the Association for lacrimal and salivary glands, and represents one of the leading causes Research in Vision and Ophthalmology Fort Lauderdale, of dry eye and mouth in the world. The purpose of this Florida, USA April 29-May 04, 2001 DOCUMENT TYPE: study was to determine whether the profound, gender-related Conference LANGUAGE: English differences observed in the magnitude of exocrine gland inflammation in SUMMARY LANGUAGE: English Sjogren's L4 ANSWER 12 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL syndrome may also be found in tissues of mouse models of this ABSTRACTS INC. disorder. ACCESSION NUMBER: 2001:78070 BIOSIS Lacrimal and submandibular glands were obtained from adult MRL/lpr, MRL+/+

(MRL+), NZB/NZW F1 (F1), C3H/lpr, C3H/gld (gld), C57BL/6lpr/lpr (B6/lpr;

with (bcl-2+/lpr) or without (bcl-2-/lpr) bcl-2 transgene insertion) and nonobese diabetic (NOD) mice after the onset of autoimmune

disease, and processed for microscopy and image analysis. Our results showed that: (1)

the extent of inflammation was significantly greater in lacrimal

female MRL/lpr, MRL+, F1, C3H/lpr and gld mice, and salivary glands of

female MRL+, F1 and gld mice, relative to those of males; (2) the severity

of inflammation in NOD mice showed a tissue-specific pattern:

DOCUMENT NUMBER: PREV200100078070

TITLE: Improvement of corneal barrier function by the P2Y2 agonist

INS365 in a rat dry eye model.

AUTHOR(S): Fujihara, Tsutomu (1); Murakami, Tadahiro; Fujita, Hiromi;

Nakamura, Masatsugu; Nakata, Katsuhiko

CORPORATE SOURCE: (1) Nara Research and Development Center, Santen

Pharmaceutical Co., Ltd., 8916-16 Takayama-cho, Ikomashi.

Nara, 630-0101: fujiharat@santen.co.jp Japan

SOURCE: IOVS, (January, 2001) Vol. 42, No. 1, pp. 96-100. print.

DOCUMENT TYPE: Article LANGUAGE: English
SUMMARY LANGUAGE: English

AB Purpose. Because purinoceptor P2Y2 receptor agonists elicit increases in

net Cl, fluid transport, and glycoprotein release onto the ocular

they are candidates for treatment of **dry** eye syndrome.

Accordingly, the effects of such an agonist INS365 on these

were characterized in a rat dry eye model. Methods. An SD rat dry eye model was used in which exorbital

lacrimal gland extirpation decreased the Schirmer test score by at

50%. After 8 weeks, when significant increases occurred in corneal epithelial permeability, INS365-containing eye drops were applied six

times daily for the next 4 weeks at concentrations from 0.03% to 3.0%.

Corneal barrier function was evaluated based on measurements with

modified anterior fluorometer of fluorescein penetrance at 1, 2, and 4 weeks after initial application. After INS365 application, the periodic acid-Schiff reagent (PAS)-stained area was evaluated in histologic sections of the tarsal and bulbar conjunctiva. Results. Ten minutes after

INS365 eye drop application at doses of either 3.0% or 8.5%, a 1.5-fold

transient increase in tear fluid secretion occurred in both the control and dry eye model animals. These transient increases nearly returned to baseline after 60 minutes. Furthermore, after 5 minutes, 1.0% INS365 was sufficient to cause a maximal transient decrease

in the PAS-stained area of more than 30%, which thereafter recovered

toward the initial level. Beginning at 2 weeks and continuing for an additional 2 weeks, maximal declines in dye penetrance of approximately

50% occurred with doses of INS365 as low as 1%. Such improvement in

corneal epithelial resistance was accompanied by complete restoration of

the PAS-stained area to the level seen in the control animal. Conclusions.

In a rat dry eye model, the P2Y2 agonist INS365 was found to improve surface health, based on increases in tear fluid secretion, corneal epithelial resistance, and release of glycoprotein-containing moieties from goblet cells. These effects suggest

that INS365 is a potential therapeutic agent for use in the treatment of dry eye syndrome.

L4 ANSWER 11 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2001:287465 BIOSIS DOCUMENT NUMBER: PREV200100287465

TITLE: Corneal protection by the ocular mucin secretagogue 15(S)-HETE in a rabbit **dry** eye model.

AUTHOR(S): Gamache, D. A. (1); Wei, Z.-Y. (1); Weimer, L. K. (1);

Miller, S. T. (1); Spellman, J. M. (1); Yanni, J. M. (1) CORPORATE SOURCE: (1) Pharmaceutical Products Research, Alcon Research, Ltd.,

Fort Worth, TX, 76134 USA

SOURCE: IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S177. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

DOCUMENT TYPE: Conference LANGUAGE: English SUMMARY LANGUAGE: English

L4 ANSWER 9 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2001:321201 BIOSIS DOCUMENT NUMBER: PREV200100321201

TITLE: A new animal model of dry

eyes.

AUTHOR(S): Nemet, A. (1); Rosner, M. (1); Belkin, M. (1) CORPORATE SOURCE: (1) Goldschleger Eye Research Institute, Tel-Aviv

University, Sheba Medical Center, Tel Hashomer Israel SOURCE: IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S262. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

DOCUMENT TYPE: Conference LANGUAGE: English SUMMARY LANGUAGE: English

L4 ANSWER 7 OF 43 BIOSIS COPYRIGHT 2002 BIOLOGICAL

ABSTRACTS INC.

ACCESSION NUMBER: 2001:321206 BIOSIS DOCUMENT NUMBER: PREV200100321206

TITLE: A new dry-eye animal model using GVH reactions.

AUTHOR(S): Uno, T. (1); Urashima, H. (1); Ohashi, Y. (1) CORPORATE SOURCE: (1) Department of Ophthalmology, Ehime University,

Matsuyama, Ehime Japan

SOURCE: IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S263. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision and Ophthalmology Fort Lauderdale,

Florida, USA April 29-May 04, 2001 DOCUMENT TYPE: Conference LANGUAGE: English

SUMMARY LANGUAGE: English

=> d his

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

L1 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

L2 251329 S ANIMAL MODEL

L3 49 S L1 AND L2

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

=> log hold

COST IN U.S. DOLLARS SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST 97.79 98.00

SESSION WILL BE HELD FOR 60 MINUTES STN INTERNATIONAL SESSION SUSPENDED AT 10:42:42 ON 01 SEP 2002

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1636DXS

PASSWORD:

***** * RECONNECTED TO STN INTERNATIONAL ****** SESSION RESUMED IN FILE 'MEDLINE, BIOSIS, CAPLUS' AT 10:45:20 ON 01 SEP 2002

FILE 'MEDLINE' ENTERED AT 10:45:20 ON 01 SEP 2002 FILE 'BIOSIS' ENTERED AT 10:45:20 ON 01 SEP 2002 COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R) FILE 'CAPLUS' ENTERED AT 10:45:20 ON 01 SEP 2002 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS

SINCE FILE TOTAL ENTRY SESSION

FULL ESTIMATED COST

97.79 98.00

=> d his

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

L1 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

L2 251329 S ANIMAL MODEL

L3 49 S L1 AND L2

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

=> s comea? epitheli?

L5 11300 CORNEA? EPITHELI?

=> s damage

L6 631428 DAMAGE

=> s damag?

L7 708684 DAMAG?

=> s dessicat? or dry

L8 565797 DESSICAT? OR DRY

=> s 15(s)16

L9 486 L5(S) L6

=> s 15(s)18

L10 142 L5(S) L8

=> s 12 and 110

L11 5 L2 AND L10

=> dup rem 111

PROCESSING COMPLETED FOR L11

L12 5 DUP REM L11 (0 DUPLICATES REMOVED)

=> d ti so 1-5

L12 ANSWER 1 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Improvement of corneal barrier function by the P2Y2 agonist INS365 in a

rat dry eye model.

SO IOVS, (January, 2001) Vol. 42, No. 1, pp. 96-100. print.

L12 ANSWER 2 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The study of TEM and SEM in the rabbit cornea after excimer laser photorefractive keratectomy.

SO Xi'an Yike Daxue Xuebao, (Apr., 2000) Vol. 21, No. 2, pp. 134-136. print.

ISSN: 0258-0659.

L12 ANSWER 3 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term dry eye model.

SO Journal of Ocular Pharmacology and Therapeutics, (April, 1998) Vol. 14.

No. 2, pp. 99-107.

ISSN: 1080-7683.

L12 ANSWER 4 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Gefarnate stimulates secretion of mucin-like glycoproteins by corneal

epithelium in vitro and protects corneal epithelium from desiccation

in

vivo.

SO Experimental Eye Research, (1997) Vol. 65, No. 4, pp. 569-574. ISSN: 0014-4835.

L12 ANSWER 5 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The effects of artificial tear solutions on wound healing in full thickness corneal incisions.

SO Acta Physiologica Hungarica, (1997) Vol. 85, No. 3, pp. 251-258. ISSN: 0231-424X.

=> d ibib ab 4,3

L12 ANSWER 4 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL

ABSTRACTS INC.

ACCESSION NUMBER: 1997:508567 BIOSIS DOCUMENT NUMBER: PREV199799807770

TITLE: Gefarnate stimulates secretion of mucin-like

glycoproteins

by corneal epithelium in vitro and protects corneal epithelium from desiccation in vivo.

AUTHOR(S): Nakamura, Masatsugu (1); Endo, Ken-Ichi; Nakata, Katsuhiko;

Hamano, Takashi

CORPORATE SOURCE: (1) Ophthalmic Lab., Nara Res. Dev. Cent.,

Santen

Pharmaceuical Co. Ltd., 8916-16 Takayama-cho, Ikoma-

shi,

Nara 630-01 Japan

SOURCE: Experimental Eye Research, (1997) Vol. 65, No. 4,

pp.

569-574.

ISSN: 0014-4835.

DOCUMENT TYPE: Article LANGUAGE: English

AB The effect of drugs for gastritis and gastric ulcer (ecabet sodium, gefarnate, teprenone, and troxipide) on the secretion of mucin-like glycoproteins from rat cornea were investigated in vitro and on a short-term, rabbit dry eye model in vivo. For the studies in

vitro, cultured rat comea sections (3 mm diameter) were incubated with

radiolabeled sodium sulfate, rinsed, and then incubated for 30 min in the

presence of one of the drugs. The culture media were reacted with Dolichos

biflorus agglutinate (DBA)-lectin, and the radioactivity of DBA-bound

mucin-like glycoproteins was measured. A cytotoxicity assay confirmed that

mucin-like glycoproteins had not leaked from damaged cells. For studies in vivo, eye drop vehicle or drops containing gefarnate were instilled in

the eyes of nine anesthetized rabbits, and then the eyes were kept open

with specula for two hours. These rabbits and two control rabbits not subjected

to ocular drying were killed, and their eyes were enucleated and stained

with methylene blue. Corneal epithelial damage from desiccation was evaluated based on the extent of methylene blue taining.

Among the four kinds of drugs for gastritis and gastric ulcers, only gefarnate significantly increased the mucin-like glycoprotein secretion

from cultured rat corneas in vitro: this stimulatory effect of gefarnate was dose-dependent. In vivo, the instillation of gefarnate reduced corneal epithelial damage from desiccation in a

dose-dependent fashion. These results suggest that gefarnate reduces desiccation of corneal epithelium, perhaps by

stimulating secretion of mucin-like glycoproteins from corneal epithelium.

L12 ANSWER 3 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1998:261667 BIOSIS

DOCUMENT NUMBER: PREV199800261667
TITLE: Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term

dry eye model.

AUTHOR(S): Fujihara, Tsutomu (1); Nagano, Takashi; Nakamura,

Masatsugu; Shirasawa, Eiichi

CORPORATE SOURCE: (1) Santen Pharmaceutical Co., Ltd. Nara Res. Develop.

Cent., 8916-16, Takayama-cho, Ikoma-shi, Nara, 630-01

Japan

SOURCE: Journal of Ocular Pharmacology and Therapeutics,

(April,

1998) Vol. 14, No. 2, pp. 99-107.

ISSN: 1080-7683.

DOCUMENT TYPE: Article LANGUAGE: English

AB Human tear fluid contains lactoferrin at the highest concentration.

patients with dry eye such as Sjogren's syndrome, the concentration of lactoferrin in the tears is approximately half the ormal

value. The present study utilizes a short-term rabbit dry eye model to evaluate if lactoferrin containing eye drops can reverse any

the damage produced by blockage of blinking with an ocular speculum.

Damage was evaluated based on the extent of methylene blue staining in

histological sections. After 3 h of desiccation, the amount of extractable dye recovered following sacrifice increased by more than 4-fold in the vehicle-treated eyes. However, in those rabbits treated with

1% lactoferrin, dye recovery was only 40% of the value in the vehicle-treated eyes. Between 1-3 h and over a concentration range from

0.01 to 1% lactoferrin, the decreases in staining were both time and concentration dependent. Alternatively, if 1% lactoferrin was applied during the desiccation period, there was partial restoration of corneal epithelial integrity. These results suggest that lactoferrin may be of therapeutic value in decreasing the loss of corneal epithelial integrity in dry eye.

=> d his

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

L1 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

L2 251329 S ANIMAL MODEL

L3 49 S L1 AND L2

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

L5 11300 S CORNEA? EPITHELI?

L6 631428 S DAMAGE

L7 708684 S DAMAG?

L8 565797 S DESSICAT? OR DRY

L9 486 S L5(S)L6

L10 142 S L5(S)L8

L11 5 S L2 AND L10

L12 5 DUP REM L11 (0 DUPLICATES REMOVED)

=> s 15(S)17

L13 586 L5(S) L7

=> s 113 and 18

L14 32 L13 AND L8

=> dup rem 114

PROCESSING COMPLETED FOR L14

L15 21 DUP REM L14 (11 DUPLICATES REMOVED)

=> d ti so 1-21

L15 ANSWER 1 OF 21 MEDLINE

TI Canaliculitis caused by Actinomyces in a case of dry eye with punctal plug occlusion.

SO NIPPON GANKA GAKKAI ZASSHI. ACTA SOCIETATIS OPHTHALMOLOGICAE JAPONICAE,

(2002 Jul) 106 (7) 416-9.

Journal code: 7505716. ISSN: 0029-0203.

L15 ANSWER 2 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The effect of tear film break-up on higher-order aberrations measured with

wavefront sensor.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S900. print. Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

L15 ANSWER 3 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI A new dry-eye animal model using GVH reactions.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S263. print.

Meeting Info.: Annual Meeting of the Association for Research in Vision

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

L15 ANSWER 4 OF 21 MEDLINE DUPLICATE

TI Bilateral neurotrophic keratopathy complicating Vidian neurectomy. SO AMERICAN JOURNAL OF OPHTHALMOLOGY, (2001 Jul) 132 (1) 106-8.

Journal code: 0370500. ISSN: 0002-9394.

L15 ANSWER 5 OF 21 MEDLINE

TI Interactions between ocular surface fluid and cornea related to contact

lenses.

vivo.

SO EUROPEAN JOURNAL OF OPHTHALMOLOGY, (2001 Apr-Jun) 11 (2) 105-15. Ref: 53

Journal code: 9110772. ISSN: 1120-6721.

L15 ANSWER 6 OF 21 CAPLUS COPYRIGHT 2002 ACS

TI Dry eye treatment with eye drops that stimulate mucin production SO Advances in Experimental Medicine and Biology (1998), 438(Lacrimal Gland,

Tear Film, and Dry Eye Syndromes 2), 965-968 CODEN: AEMBAP; ISSN: 0065-2598

L15 ANSWER 7 OF 21 MEDLINE

DUPLICATE 2

TI Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term dry eye model.

SO JOURNAL OF OCULAR PHARMACOLOGY AND

THERAPEUTICS, (1998 Apr) 14 (2) 99-107.

Journal code: 9511091. ISSN: 1080-7683.

L15 ANSWER 8 OF 21 MEDLINE

DUPLICATE 3

TI Gefarnate stimulates secretion of mucin-like glycoproteins by corneal

epithelium in vitro and protects corneal epithelium from desiccation in

SO EXPERIMENTAL EYE RESEARCH, (1997 Oct) 65 (4) 569-74. Journal code: 0370707. ISSN: 0014-4835.

L15 ANSWER 9 OF 21 MEDLINE

DUPLICATE 4

TI Evaluation of conjunctival epithelial damage in dry eye.

SO NIPPON GANKA GAKKAI ZASSHI. ACTA SOCIETATIS OPHTHALMOLOGICAE JAPONICAE,

(1997 Jan) 101 (1) 52-6.

Journal code: 7505716. ISSN: 0029-0203.

L15 ANSWER 10 OF 21 MEDLINE

DUPLICATE 5

TI Establishment of a rabbit short-term dry eye model. SO JOURNAL OF OCULAR PHARMACOLOGY AND

THERAPEUTICS, (1995 Winter) 11 (4)

Journal code: 9511091. ISSN: 1080-7683.

- L15 ANSWER 11 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI Effects of preservative-free artificial tear solutions on corneal epithelial structure and function.
- SO Archives of Ophthalmology, (1995) Vol. 113, No. 3, pp. 371-378. ISSN: 0003-9950.
- L15 ANSWER 12 OF 21 MEDLINE
- TI Artificial tear composition and promotion of recovery of the damaged corneal epithelium.
- SO CORNEA, (1993 Mar) 12 (2) 115-20. Journal code: 8216186. ISSN: 0277-3740.
- L15 ANSWER 13 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE

TI SURFACE PROPERTIES OF NORMAL AND DAMAGED CORNEAL

EPITHELIA.

- SO J DISPERSION SCI TECHNOL, (1992) 13 (4), 459-478. CODEN: JDTEDS. ISSN: 0193-2691.
- L15 ANSWER 14 OF 21 MEDLINE **DUPLICATE 7** TI Histochemical changes in the rabbit cornea and plasmin activity in the

tear fluid during contact lens wear. Favourable influence of protease inhibitors (aprotinin, PC5, elastatinal).

- SO HISTOCHEMISTRY, (1992) 97 (1) 69-76. Journal code: 0411300. ISSN: 0301-5564.
- L15 ANSWER 15 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI TOXICITY OF EYEDROPS WITH BENZALKONIUM CHLORIDE ASSESSMENT VIA TEAR

LACTATE DEHYDROGENASE ACTIVITY.

- SO FOLIA OPHTHALMOL JPN, (1991) 42 (5 PART 1), 780-783. CODEN: NGKYA3. ISSN: 0015-5667.
- **DUPLICATE 8** L15 ANSWER 16 OF 21 MEDLINE
- TI Quantitative evaluation of the corneal epithelial barrier: effect of artificial tears and preservatives.
- SO CURRENT EYE RESEARCH, (1991 Jul) 10 (7) 645-56. Journal code: 8104312. ISSN: 0271-3683.
- L15 ANSWER 17 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
- TI DRY EYE FOLLOWING TOPICAL BETA-BLOCKERS.
- SO ANN OTTALMOL CLIN OCUL, (1990) 116 (10), 989-992. CODEN: AOCOAG. ISSN: 0003-4665.
- L15 ANSWER 18 OF 21 MEDLINE

DUPLICATE 9

- TI Influence of artificial tears on corneal epithelium in dry-eye syndrome.
- SO GRAEFES ARCHIVE FOR CLINICAL AND EXPERIMENTAL OPHTHALMOLOGY, (1989) 227

(2) 139-41.

Journal code: 8205248. ISSN: 0721-832X.

- **DUPLICATE 10** L15 ANSWER 19 OF 21 MEDLINE
- TI Dry mass and water content in the corneal epithelium and superficial stroma during healing of corneal alkali wounds.
- SO EXPERIMENTAL EYE RESEARCH, (1988 May) 46 (5) 705-15. Journal code: 0370707. ISSN: 0014-4835.

- L15 ANSWER 20 OF 21 MEDLINE
- TI Evelid secretions and the prevention and production of disease.
- SO Eye, (1988) 2 (Pt 2) 164-71. Ref: 66

Journal code: 8703986. ISSN: 0950-222X.

- L15 ANSWER 21 OF 21 MEDLINE
- TI [The lacrimal film, structure and stability].

Le film lacrymal, structure et stabilite.

SO JOURNAL FRANCAIS D OPHTALMOLOGIE, (1983) 6 (12) 963-9.

Journal code: 7804128. ISSN: 0181-5512.

=> d ibib ab 13,12,10,7

L15 ANSWER 13 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE

ACCESSION NUMBER: 1992:438754 BIOSIS

DOCUMENT NUMBER: BA94:90879 SURFACE PROPERTIES OF NORMAL AND TITLE:

DAMAGED

CORNEAL EPITHELIA.

AUTHOR(S): SHARMA A

CORPORÀTE SOURCE: DEP. OF CHEMICAL ENGINEERING, INDIAN INSTITUTE OF TECHNOL.

AT KANPUR, KANPUR 20816, INDIA.

SOURCE: J DISPERSION SCI TECHNOL, (1992) 13 (4), 459-478.

CODEN: JDTEDS. ISSN: 0193-2691.

FILE SEGMENT: BA; OLD LANGUAGE: **English**

AB Surface properties of glycoprotein strands (glycocalyx) anchored on a

variety of cell surface control many cell events such as cell adhesion, fusion, wetting, dispersion and polymer mediated flocculation. The apolar

and polar surface properties of the corneal mucus, glycocalix bearing

normal corneal epithelial cells, and damaged

corneal cells were determined by contact angle goniometry on freshly enucleated rabbit corneas. The corneal mucus, as well as the cell glycocalyx, display intense electron donor or proton acceptor (basic) type

of monopolarity, because of which both have negative interfacial tensions

against water, and both are almost as hydrophilic as water itself. These

findings show that the mucus covering of the cornea is neither necesssary

for the corneal wetting by tears, nor is its loss sufficient for explanation of the corneal nonwettability. However, damage and desiccation of the epithelial cell and associated glycocalix resulted in substantial loss of the polar surface properties, and in diminished hydrophilicity and wettability. This observation supports the hypothesis

that cell damage and glycocalyx abnormalities may lead to

L15 ANSWER 12 OF 21 MEDLINE

ACCESSION NUMBER: 93272522 MEDLINE DOCUMENT NUMBER: 93272522 PubMed ID: 8500317

Artificial tear composition and promotion of recovery of TITLE: the damaged corneal epithelium

Lopez Bernal D; Ubels J L AUTHOR:

CORPORATE SOURCE: Department of Ophthalmology, Medical College of Wisconsin,

Milwaukee.

CONTRACT NUMBER: P30 EY01931 (NEI)

CORNEA, (1993 Mar) 12 (2) 115-20. SOURCE:

Journal code: 8216186. ISSN: 0277-3740.

United States PUB. COUNTRY:

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English FILE SEGMENT: Priority Journals

199306 ENTRY MONTH:

Entered STN: 19930716 ENTRY DATE: Last Updated on STN: 19930716 Entered Medline: 19930630

AB In severe dry eye syndromes the corneal

epithelium is compromised with development of punctate erosions and increased permeability. In the present study the ability of

tear solutions to promote recovery of the corneal epithelial barrier was determined by measurement of corneal uptake of 5,6 carboxyfluorescein (CF). Corneas of anesthetized rabbits were exposed to 0.01% benzalkonium for 5 min to increase epithelial permeability. The cornea was then exposed to an artificial tear

for 1.5 h followed by measurement of CF uptake. During exposure to

commercial isotonic, nonpreserved solutions and a solution preserved with

polyquaternium-1, CF uptake decreased significantly but did not return to

control. No recovery of the epithelial barrier occurred during exposure of

corneas to nonpreserved hypotonic solutions. During exposure to an experimental tear solution with an electrolyte composition similar to human tears, buffered with bicarbonate, CF uptake returned to control

levels. Bicarbonate is an essential component of this solution because the

same formula buffered with borate or without buffer was ineffective

promoting recovery of the damaged corneal epithelium.

DUPLICATE 5 L15 ANSWER 10 OF 21 MEDLINE ACCESSION NUMBER: 96153755 MEDLINE

DOCUMENT NUMBER: 96153755 PubMed ID: 8574813 Establishment of a rabbit short-term dry eye TITLE:

Fujihara T; Nagano T; Nakamura M; Shirasawa E AUTHOR: CORPORATE SOURCE: Santen Pharmaceutical Co., Ltd., Ophthalmic Research

Laboratories, Osaka, Japan.

JOURNAL OF OCULAR PHARMACOLOGY AND SOURCE: THERAPEUTICS, (1995

Winter) 11 (4) 503-8.

Journal code: 9511091. ISSN: 1080-7683.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals ENTRY MONTH: 199603

Entered STN: 19960321 ENTRY DATE: Last Updated on STN: 19960321

Entered Medline: 19960311

AB The dry eye syndrome is a chronic disease which can become a serious threat to useful vision. However, there is only a limited understanding regarding effective treatment or prevention of this disease.

Establishing an effective mode of treatment requires the use of a satisfactory animal dry eye model. Ideally, such a model should rapidly determine the effectiveness of agents that inhibit the damaging effects of this syndrome. This paper presents a short-term dry eye model using rabbits, which combines mechanical prevention of blinking and methylene blue staining. This

is not intended to be a precise representation of the dry eye syndrome, since this disorder has recently become recognized to involve a

primary pathological process of the corneal and conjunctival epithelium.

However, by using this model, clinical signs of dry eye can be observed after a few hours in the form of acute desiccation. Corneal damage can easily be evaluated both qualitatively by methylene blue staining scores, and quantitatively by chronic assay. Visually observed corneal epithelial thinning was confirmed by scanning electron microscopy (SEM) to be due to loss of epithelial integrity. Using a 3% chondroitin sulfate solution, an already proven effective agent for dry eye, this model effectively demonstrated an 80% inhibition in the development of methylene blue positive

after a period of only 2 hours. This short term dry eye model is valuable in primarily screening the efficacy of potential therapeutic agents in the prevention and treatment of dry eye.

L15 ANSWER 7 OF 21 MEDLINE ACCESSION NUMBER: 1998232247 MEDLINE **DUPLICATE 2** DOCUMENT NUMBER: 98232247 PubMed ID: 9572535 Lactoferrin suppresses loss of corneal epithelial TITLE: integrity

in a rabbit short-term dry eye model.

Fujihara T; Nagano T; Nakamura M; Shirasawa E AUTHOR: CORPORATE SOURCE: Santen Pharmaceutical Co., Ltd., Nara Research and

Development Center, Japan.

JOURNAL OF OCULAR PHARMACOLOGY AND SOURCE:

THERAPEUTICS, (1998 Apr)

14 (2) 99-107.

Journal code: 9511091. ISSN: 1080-7683.

United States PUB. COUNTRY:

Journal; Article; (JOURNAL ARTICLE) DOCUMENT TYPE:

LANGUAGE: English FILE SEGMENT: Priority Journals

ENTRY MONTH: 199806 Entered STN: 19980708 ENTRY DATE:

Last Updated on STN: 19980708 Entered Medline: 19980622

AB Human tear fluid contains lactoferrin at the highest concentration. In

patients with dry eye such as Sjogren's syndrome, the concentration of lactoferrin in the tears is approximately half the

value. The present study utilizes a short-term rabbit dry eye model to evaluate if lactoferrin containing eye drops can reverse any

the damage produced by blockage of blinking with an ocular speculum. Damage was evaluated based on the extent of methylene blue staining in histological sections. After 3 h of desiccation, the amount of extractable dye recovered following sacrifice increased by more

than 4-fold in the vehicle-treated eyes. However, in those rabbits treated

with 1% lactoferrin, dye recovery was only 40% of the value in the vehicle-treated eyes. Between 1-3 h and over a concentration range

0.01 to 1% lactoferrin, the decreases in staining were both time and concentration dependent. Alternatively, if 1% lactoferrin was applied during the desiccation period, there was partial restoration of corneal epithelial integrity. These results suggest that lactoferrin may be of therapeutic value in decreasing the loss of corneal epithelial integrity in dry eye.

=> D HIS

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON

4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR L1 KERATOCONJUCTIVITIS SIC

251329 S ANIMAL MODEL L2

L3 49 S L1 AND L2

43 DUP REM L3 (6 DUPLICATES REMOVED) L4

11300 S CORNEA? EPITHELI? L5

631428 S DAMAGE L6

708684 S DAMAG? 1.7

565797 S DESSICAT? OR DRY Journal code: 8505423. ISSN: 0814-9763. L8 486 S L5(S)L6 L9 142 S L5(S)L8 L18 ANSWER 6 OF 9 MEDLINE **DUPLICATE 2** L10 5 S L2 AND L10 TI Corneal epithelial changes in diabetic rats. L11 5 DUP REM L11 (0 DUPLICATES REMOVED) SO OPHTHALMIC RESEARCH, (1982) 14 (4) 269-78. L12 Journal code: 0267442. ISSN: 0030-3747. L13 586 S L5(S)L7 L14 32 S L13 AND L8 L18 ANSWER 7 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL 21 DUP REM L14 (11 DUPLICATES REMOVED) L15 ABSTRACTS INC. TI CORNEAL EPITHELIUM IN DIABETES. => OSMOT? OR OSMOS? OSMOT? IS NOT A RECOGNIZED COMMAND SO ACTA SOC OPHTHALMOL JPN, (1981) 85 (9), 1216-1225. CODEN: NGZAA6. ISSN: 0029-0203. The previous command name entered was not recognized by the system. For a list of commands available to you in the current file, enter "HELP COMMANDS" at an arrow prompt (=>). L18 ANSWER 8 OF 9 MEDLINE **DUPLICATE 3** TI Diabetes mellitus and the rabbit corneal epithelium. => S OSMOT? OR OSMOS? SO INVESTIGATIVE OPHTHALMOLOGY AND VISUAL L16 130776 OSMOT? OR OSMOS? SCIENCE, (1981 Aug) 21 (2) 317-21. Journal code: 7703701. ISSN: 0146-0404. => S L9 AND L16 L18 ANSWER 9 OF 9 MEDLINE 14 L9 AND L16 **DUPLICATE 4** TI Factors related to corneal epithelial complications after closed vitrectomy in diabetics. => DUP REM L17 PROCESSING COMPLETED FOR L17 SO ARCHIVES OF OPHTHALMOLOGY, (1979 Jun) 97 (6) 1076-8. Journal code: 7706534. ISSN: 0003-9950. 9 DUP REM L17 (5 DUPLICATES REMOVED) L18 => D TI SO 1-9 => D HIS L18 ANSWER 1 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE 1 (FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002) TI Aqueous humour and crystalline lens changes associated with FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON ultraviolet radiation or mechanical damage to corneal 01 SEP 2002 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR epithelium in freshwater rainbow trout eyes. L1 SO Journal of Photochemistry and Photobiology B Biology, (Nov., KERATOCONJUCTIVITIS SIC 1997) Vol. 251329 S ANIMAL MODEL L2 41, No. 1-2, pp. 165-172. L3 49 S L1 AND L2 ISSN: 1011-1344. L4 43 DUP REM L3 (6 DUPLICATES REMOVED) 1.5 11300 S CORNEA? EPITHELI? L18 ANSWER 2 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL L6 631428 S DAMAGE ABSTRACTS INC. L7 708684 S DAMAG? TI DIFFERENTIAL EFFECT OF PH ON THE OSMOTIC L8 565797 S DESSICAT? OR DRY TOLERANCE OF RABBIT L9 486 S L5(S)L6 CORNEAL EPITHELIUM IN TISSUE CULTURE. 142 S L5(S)L8 L10 SO J TOXICOL CUTANEOUS OCUL TOXICOL, (1991) 10 (1-2), L11 5 S L2 AND L10 5 DUP REM L11 (0 DUPLICATES REMOVED) L12 CODEN: JTOTDO. ISSN: 0731-3829. 586 S L5(S)L7 L13 L14 32 S L13 AND L8 21 DUP REM L14 (11 DUPLICATES REMOVED) L18 ANSWER 3 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL L15 130776 S OSMOT? OR OSMOS? ABSTRACTS INC. L16 TI OSMOTIC TOLERANCE OF RABBIT CORNEAL L17 14 S L9 AND L16 EPITHELIUM IN TISSUE 9 DUP REM L17 (5 DUPLICATES REMOVED) L18 CULTURE. SO J TOXICOL CUTANEOUS OCUL TOXICOL, (1991) 10 (1-2), => LOG HOLD 59-78. COST IN U.S. DOLLARS SINCE FILE TOTAL CODEN: JTOTDO. ISSN: 0731-3829. ENTRY SESSION **FULL ESTIMATED COST** 163.15 163.36 L18 ANSWER 4 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC. SESSION WILL BE HELD FOR 60 MINUTES TI ACUTE AND PROTRACTED DAMAGE TO THE CORNEAL STN INTERNATIONAL SESSION SUSPENDED AT 11:04:19 ON 01 EPITHELIUM IN-VITRO IS A INDICATION OF SEP 2002 BIOCOMPATIBILITY OF PHARMACEUTICAL VEHICLES. Connecting via Winsock to STN SO ANNUAL SPRING MEETING OF THE ASSOCIATION FOR RESEARCH IN VISION AND OPHTHALMOLOGY, SARASOTA, FLORIDA, USA, MAY 1-6, 1988. INVEST OPHTHALMOL Welcome to STN International! Enter x:x VISUAL SCI. (1988) 29 (ABSTR ISSUE), 193. CODEN: IOVSDA. ÍSSN: 0146-0404. LOGINID:SSSPTA1636DXS

L18 ANSWER 5 OF 9 MEDLINE PASSWORD: TI Corneal endothelial changes in superficial epithelial keratopathy. * * * * * * RECONNECTED TO STN INTERNATIONAL * * * * * SO AUSTRALIAN AND NEW ZEALAND JOURNAL OF SESSION RESUMED IN FILE 'MEDLINE, BIOSIS, CAPLUS' AT OPHTHALMOLOGY, (1986 May) 14 (2) 11:09:48 ON 01 SEP 2002 FILE 'MEDLINE' ENTERED AT 11:09:48 ON 01 SEP 2002

171-5.

FILE 'BIOSIS' ENTERED AT 11:09:48 ON 01 SEP 2002 COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R) FILE 'CAPLUS' ENTERED AT 11:09:48 ON 01 SEP 2002 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS

SINCE FILE TOTAL

SO J CATHOL MED COLL, (1992) 45 (2), 609-621. CODEN: KTUNAA. ISSN: 0368-7015.

OF SUPERFICIAL CONJUNCTIVAL EPITHELIUM IN

ENTRY SESSION

FULL ESTIMATED COST

163.15 163.36

=> D HIS

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

L1 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

L2 251329 S ANIMAL MODEL

L3 49 S L1 AND L2

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

L5 11300 S CORNEA? EPITHELI?

L6 631428 S DAMAGE

L7 708684 S DAMAG?

L8 565797 S DESSICAT? OR DRY

L9 486 S L5(S)L6

L10 142 S L5(S)L8

L11 5 S L2 AND L10

L12 5 DUP REM L11 (0 DUPLICATES REMOVED)

L13 586 S L5(S)L7

L14 32 S L13 AND L8

L15 21 DUP REM L14 (11 DUPLICATES REMOVED)

L16 130776 S OSMOT? OR OSMOS?

L17 14 S L9 AND L16

L18 9 DUP REM L17 (5 DUPLICATES REMOVED)

=> S HYPERTONIC

L19 29959 HYPERTONIC

=> S L1 AND L19

L20 8 L1 AND L19

=> DUP REM L20

PROCESSING COMPLETED FOR L20

L21 6 DUP REM L20 (2 DUPLICATES REMOVED)

=> D TI SO 1-6

L21 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2002 ACS

TI Adaptation by corneal epithelial cells to chronic hypertonic stress depends on upregulation of Na:K:2Cl cotransporter gene and protein

expression and ion transport activity

SO Journal of Membrane Biology (2000), 177(1), 41-50 CODEN: JMBBBO; ISSN: 0022-2631

CODEN: JMBBBO; 188N: 0022-203

L21 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2002 ACS

TI Effects of chronic hypertonic stress on regulatory volume increase and Na-K-2Cl cotransporter expression in cultured corneal epithelial cells

SO Advances in Experimental Medicine and Biology (1998), 438(Lacrimal Gland,

Tear Film, and Dry Eye Syndromes 2), 637-642 CODEN: AEMBAP; ISSN: 0065-2598

L21 ANSWER 3 OF 6 MEDLINE DUPLICATE 1
TI Human basic tear fluid osmolality. II. Importance of processing

SO ACTA OPHTHALMOLOGICA, (1993 Aug) 71 (4) 530-8. Journal code: 0370347. ISSN: 0001-639X.

L21 ANSWER 4 OF 6 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI THE EFFECTS OF HYPERTONIC SOLUTIONS ON ULTRASTRUCTURAL CHANGES

L21 ANSWER 5 OF 6 MEDLINE

DUPLICATE 2

TI Tear osmotic differences across the ocular surface.

SO GRAEFES ARCHIVE FOR CLINICAL AND EXPERIMENTAL OPHTHALMOLOGY, (1986) 224

(6) 583-6.

RABBITS.

Journal code: 8205248. ISSN: 0721-832X.

L21 ANSWER 6 OF 6 MEDLINE

TI New aspects of contact lenses in ophthalmology.

SO ADVANCES IN OPHTHALMOLOGY, (1976) 32 2-34.

Journal code: 0263212. ISSN: 0065-3004.

=> D IBIB AB 1, 4

L21 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:647856 CAPLUS

DOCUMENT NUMBER: 133:361170

TITLE: Adaptation by corneal epithelial cells to chronic

hypertonic stress depends on upregulation of

Na:K:2Cl cotransporter gene and protein expression and

ion transport activity

AUTHOR(S): Bildin, V. N.; Yang, H.; Crook, R. B.; Fischbarg,

J.;

Reinach, P. S.

CORPORATE SOURCE: Department of Biological Sciences,

College of

Optometry, State University of New York, New York,

NY,

10010, USA

SOURCE: Journal of Membrane Biology (2000), 177(1), 41-

50

CODEN: JMBBBO; ISSN: 0022-2631 Springer-Verlag New York Inc.

PUBLISHER: Springer-Verlag DOCUMENT TYPE: Journal

LANGUAGE: English

AB We examd, the ability of SV40-immortalized human and rabbit

epithelial cells (HCEC and RCEC, resp.) to adapt to chronic hypertonic stress. Under isotonic conditions in the presence of 50 .mu.M bumetanide, proliferation measured as 3H-thymidine

incorporation declined in RCEC and HCEC by 8 and 35%, resp. After 48 h exposure to 375

mOsm medium, RCEC proliferation fell by 19% whereas in HCEC it declined by

45%. Light scattering behavior demonstrated that both cell lines mediate

nearly complete regulatory vol. increase (RVI) responses to an acute hypertonic (375 mOsm) challenge, which in part depend on burnetanide-sensitive Na-K-2Cl cotransporter (NKCC) activity. Following

RCEC exposure for 48 h to 375 mOsm medium, the RVI response to an acute $\,$

hypertonic challenge was inhibited by 17%. However, in HCEC this response declined by 68%. During exposure to 375 mOsm medium for up to 24

h, only RCEC upregulated NKCC gene and protein expression as well as

burnetanide-sensitive 86Rb influx. These increases are consistent with the

smaller declines in RVI and proliferation capacity which occur during this

period in RCEC compared to HCEC. Therefore, adaptation by RCEC to chronic

hypertonic stress is dependent on stimulation of NKCC gene and protein expression and functional activity. On the other hand, under isotonic conditions, HCEC RVI and proliferation are more dependent on NKCC

activity than they are in RCEC.

REFERENCE COUNT: 24 THERE ARE 24 CITED

REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L21 ANSWER 4 OF $6\,$ BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1992:507957 BIOSIS DOCUMENT NUMBER: BA94:126482

TITLE: THE EFFECTS OF HYPERTONIC SOLUTIONS ON ULTRASTRUCTURAL CHANGES OF SUPERFICIAL

CONJUNCTIVAL

EPITHELIUM IN RABBITS.

AUTHOR(S): KIM J K; RHEE S W

CORPORATE SOURCE: DEP. OPHTHALMOLOGY, CATHOLIC UNIV. MED. COLL., SEOUL,

KOREA.

SOURCE: J CATHOL MED COLL, (1992) 45 (2), 609-621.

CODEN: KTUNAA. ISSN: 0368-7015.

FILE SEGMENT: BA; OLD LANGUAGE: Korean

AB It is well known that aside from goblet cells and the lacrimal gland there

is a third source of mucus secretion on to the surface of the conjunctiva.

Deducing from transmission electron microscopy the superficial cells

the epithelium make a mucoprotein with long chemical chains and package it

in the subsurface vesicles. This content is a particular mucoprotein, the

glycocalyx which participates in the elaboration of the cell coat and might be serving as an intermediary on the fixation of overlying

the epithelial surface. The epithelial surface is probably the most

force on maintaining the stability of the tear film and it is here that the adhesive forces must act. Epithelial cell integrity is essential for the stability of the tear film especially in dry eyes.

Elevated tear film osmolarity is a final common pathway for the development of many types of "dry eyes". Tear film

osmolarity may increase due to decreased tear secretion and/or increased

tear film evaporation. The aim of this study is to evaluate the morphologic effect of hyperosmolarity, equivalent to that seen on keratoconjunctivitis on conjunctival surface epithelium of rabbits. **Hypertonic** solutions were prepared by a commercially available balanced salt solution (BSS, Alcon), and adjusted to desired final concentrations of 300, 330, and 360 m Osm/1. New Zealand white beta.

weighing 2-3 kg were anesthetized, and a conjunctival well which was made

by elevating the lids and nictitating membrane, was filled with 1 ml of

the test solution. The conjunctival sac was submerged in a bathing solution that was exchanged for same fresh solution every hours for 3 to 6

hours. In twelve experimental trials, BSS concentrated to 330, 360 m Osm/l

was used to bathe the ocular surface for 3 and 6 hour periods, respectively. The contralateral twelve eyes bathed in isotonic BSS erved

as control. Immediately after conjunctival bathing, biopsy samples

taken from inferonasal bulbar conjunctiva. The biopsy samples were stained with ruthenium red stock solution contrasted with uranyl

and lead citrate, and the experimental specimens and their controls were

examined in a transmission electron microscope (Joel 1200 EX, Japan). And

the results were analyzed by an image analysis system (Kontron, West

Germany). The results were as follows: 1) After 3 hours of bathing in 330

m Osm/l solution, the subsurface vesicles were observed in clusters and

increased in number compared to the control group. After 6 hours of bathing in 330 m Osm/l solution, the vesicles increased in number compared

to the 3 hour group. 2) After 3 hours of bathing in 360 m Osm/l solution,

the vesicles were morphologically enlarged and increased in number compared to the control group. After 6 hours of bathing in 360 m Osm/l

solution, the vesicles were not morphologically changed compared to the

330 m Osm/l group. 3) By an image analysis system, after 3 hours of bathing in 360 m Osm/l solution, the vesicles significantly increased by

three times in total volume compared to the control group (P (0.05). After

6 hours of bathing in 330 m Osm/l solution, the vesicles significantly increased in total volume by about two times compared to the 3 hour group

(P (0.05). 4) In experimental specimens, the conjunctival epithelial cells

had widened intercellular spaces and intracellular edema, manifested by

decreased cytoplasmic density, and the microvilli towards the preocular

tear film did not appear to be reduced by increasing osmolarity, although

their shape were irregular. The above findings suggest that subsurface vesicles increase early in number and total volume in the hyperosmolar

state, at a later time the altered subsurface vesicles by the hyperosmolarity will induce the alteration of mucus content. And the conjunctival epithelium would lose the capacity to synthesize glycocalyx

and lack the specific sites of fixation for these mucoproteins.

=> D HIS

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

L1 4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

L2 251329 S ANIMAL MODEL

L3 49 S L1 AND L2

L4 43 DUP REM L3 (6 DUPLICATES REMOVED)

L5 11300 S CORNEA? EPITHELI?

L6 631428 S DAMAGE

L7 708684 S DAMAG?

L8 565797 S DESSICAT? OR DRY

L9 486 S L5(S)L6

L10 142 S L5(S)L8

L11 5 S L2 AND L10

L12 5 DUP REM L11 (0 DUPLICATES REMOVED)

L13 586 S L5(S)L7

L14 32 S L13 AND L8

L15 21 DUP REM L14 (11 DUPLICATES REMOVED)

L16 130776 S OSMOT? OR OSMOS?

L17 14 S L9 AND L16

L18 9 DUP REM L17 (5 DUPLICATES REMOVED)

L19 29959 S HYPERTONIC

L20 8 S L1 AND L19

L21 6 DUP REM L20 (2 DUPLICATES REMOVED)

=> S HYPERTONIC OR HYPEROSMO?

L22 40993 HYPERTONIC OR HYPEROSMO?

=> S L22 AND L5

L23 40 L22 AND L5

=> DUP REM L23 PROCESSING COMPLETED FOR L23 31 DUP REM L23 (9 DUPLICATES REMOVED)

=> D TI SO 1-31

L24 ANSWER 1 OF 31 MEDLINE

TI Osmosensitive taurine transporter expression and activity in human corneal epithelial cells.

SO INVESTIGATIVE OPHTHALMOLOGY AND VISUAL SCIENCE, (2002 Sep) 43 (9) 2916-22.

Journal code: 7703701. ISSN: 0146-0404.

L24 ANSWER 2 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Osmosensitive taurine transporter expression and uptake by human corneal epithelial cells.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S588. print. Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

L24 ANSWER 3 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Possible involvement of MAPKs in direct activation of Na:K:2Cl cotransporter hypertonicity-stressed rabbit and human corneal epithelial cells.

SO IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S499. print. Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04,

L24 ANSWER 4 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI MAPK involvement in recovery of paracellular resistance during

a hypertonic challenge in cultured rabbit and human corneal epithelial layers.

SO IOVS, (March 15, 2000) Vol. 41, No. 4, pp. S903. print.. Meeting Info.: Annual Meeting of the Association in Vision and Opthalmology. Fort Lauderlade, Florida, USA April 30-May 05, 2000

Association for Research in Vision and Ophthalmology

L24 ANSWER 5 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Hypertonic stress-induced activation of MAPK and apoptosis in SV40-transformed human and rabbit corneal epithelial cells.

SO IOVS, (March 15, 2000) Vol. 41, No. 4, pp. S902. print.. Meeting Info.: Annual Meeting of the Association in Vision and Opthalmology. Fort Lauderlade, Florida, USA April 30-May 05,

Association for Research in Vision and Ophthalmology

L24 ANSWER 6 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Effects of hyperosmolarity on bovine corneal epithelium cell culture. Advantages and disadvantages of using artificial tears.

SO IOVS, (March 15, 2000) Vol. 41, No. 4, pp. S67.

Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology. Fort Lauderdale, Florida, USA April 30-May 05, 2000

Association for Research in Vision and Ophthalmology

DUPLICATE 1

TI Adaptation by corneal epithelial cells to chronic hypertonic stress depends on upregulation of Na:K:2Cl cotransporter gene and protein expression and ion transport activity. SO JOURNAL OF MEMBRANE BIOLOGY, (2000 Sep 1) 177 (1) 41-50.

Journal code: 0211301. ISSN: 0022-2631.

L24 ANSWER 8 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Corneal epithelial cell adaptation to chronic

hypertonic stress is associated with upregulation of Na-K-2Cl expression and activity.

SO FASEB Journal, (March 12, 1999) Vol. 13, No. 4 PART 1, pp.

Meeting Info.: Annual Meeting of the Professional Research Scientists for

Experimental Biology 99 Washington, D.C., USA April 17-21, 1999 ISSN: 0892-6638.

L24 ANSWER 9 OF 31 MEDLINE

TI [Measurement of adhesion of the corneal lamella to the stroma in the early

postoperative period after microkeratotomy].

Mereni adheze rohovkove lamely ke stromatu po rezu

mikrokeratomem v casne

pooperacni fazi.

SO CESKA A SLOVENSKA OFTALMOLOGIE, (1999 Nov) 55 (6) 367-71.

Journal code: 9600515. ISSN: 1211-9059.

L24 ANSWER 10 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Corneal epithelial cell adaptation to chronic

hypertonic stress is associated with increase of Na-K-2Cl cotransporter expression and activity.

SO IOVS, (March 15, 1999) Vol. 40, No. 4, pp. S182.

Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology Fort Lauderdale, Florida, USA May 9-14, 1999

for Research in Vision and Opthalmology

L24 ANSWER 11 OF 31 MEDLINE

DUPLICATE 2 TI Effects of chronic hypertonic stress on regulatory volume increase and Na-K-2Cl cotransporter expression in cultured corneal

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1998) 438 637-42.

Journal code: 0121103. ISSN: 0065-2598.

L24 ANSWER 12 OF 31 MEDLINE

DUPLICATE 3

TI Reduction of water permeability by anisotonic solutions in frog corneal epithelium.

SO INVESTIGATIVE OPHTHALMOLOGY AND VISUAL SCIENCE, (1998 Feb) 39 (2) 378-84.

Journal code: 7703701. ISSN: 0146-0404.

L24 ANSWER 13 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Measurement of changes in cell volume based on fluorescence quenching.

SO American Journal of Physiology, (1997) Vol. 272, No. 4 PART 1,

C1405-C1414. ISSN: 0002-9513.

L24 ANSWER 14 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Effects of chronic hyperosmotic stress on regulatory volume recovery in cultured human and rabbit corneal epithelial

SO Investigative Ophthalmology & Visual Science, (1997) Vol. 38, No. 4 PART

1-2, pp. S1039.

Meeting Info.: Annual Meeting of the Association for Research in

and Ophthalmology, Parts 1-2 Fort Lauderdale, Florida, USA May 11-16, 1997

ISSN: 0146-0404.

L24 ANSWER 15 OF 31 MEDLINE

TI Sequential staining: the effects of sodium fluorescein, osmolarity, and pH

on human corneal epithelium.

SO OPTOMETRY AND VISION SCIENCE, (1997 Apr) 74 (4) 207-

Journal code: 8904931. ISSN: 1040-5488.

L24 ANSWER 16 OF 31 MEDLINE

TI The effect of osmolality on the shedding rate of the corneal epithelium.

SO CORNEA, (1996 May) 15 (3) 240-4. Journal code: 8216186. ISSN: 0277-3740.

L24 ANSWER 17 OF 31 MEDLINE

DUPLICATE 4

TI Evaluation of the effects of saline versus bicarbonate-containing mixed

salts solutions on rabbit corneal epithelium in vitro.

SO OPHTHALMIC AND PHYSIOLOGICAL OPTICS, (1995 Nov) 15 (6) 585-99.

Journal code: 8208839. ISSN: 0275-5408.

L24 ANSWER 18 OF 31 MEDLINE

TI Biomechanical behavior of the cornea and its response to radial keratotomy.

SO JOURNAL OF REFRACTIVE AND CORNEAL SURGERY, (1994 May-Jun) 10 (3) 343-51;

discussion 351-6.

Journal code: 9431306. ISSN: 1081-0803.

L24 ANSWER 19 OF 31 MEDLINE

TI A Na:H exchanger subtype mediates volume regulation in bovine corneal epithelial cells.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1994) 350 105-10.

Journal code: 0121103. ISSN: 0065-2598.

L24 ANSWER 20 OF 31 MEDLINE DUPLICATE 5

TI Effect of volume changes on a potassium current in rabbit corneal epithelial cells.

SO AMERICAN JOURNAL OF PHYSIOLOGY, (1993 May) 264 (5 Pt 1) C1238-45.

Journal code: 0370511. ISSN: 0002-9513.

L24 ANSWER 21 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI TOPICAL HYPEROSMOLAR SOLUTIONS IN THE REDUCTION OF CORNEAL

EDEMA.

SO CLAO (CONTACT LENS ASSOC OPHTHALMOL) J, (1987) 13 (3), 149-151.

CODEN: CLAJEU. ISSN: 0733-8902.

L24 ANSWER 22 OF 31 CAPLUS COPYRIGHT 2002 ACS

TI The effect of different solutions of glycerin on corneal thickness and cell morphology studied by specular microscopy and scanning electron

microscopy

SO Ital. J. Ophthalmol. (1987), 1(2), 67-77 CODEN: ITJOE8

L24 ANSWER 23 OF 31 CAPLUS COPYRIGHT 2002 ACS

TI Contact lens-induced edema in vitro - amelioration by lactate dehydrogenase inhibitors

SO Curr. Eye Res. (1986), 5(10), 751-8 CODEN: CEYRDM; ISSN: 0271-3683

L24 ANSWER 24 OF 31 MEDLINE

TI [Vitrectomy: measures for the prevention of intraoperative complications].

Vitrektomie: Massnahmen zur Vermeidung intraoperativer Komplikationen.

SO KLINISCHE MONATSBLATTER FUR AUGENHEILKUNDE, (1986 Apr.) 188 (4) 322-4.

Journal code: 0014133. ISSN: 0023-2165.

L24 ANSWER 25 OF 31 MEDLINE

DUPLICATE 6

TI A comparison of therapeutic bandage lenses, tarsorrhaphy, and antibiotic

and hypertonic saline on corneal epithelial wound healing.

SO ANNALS OF OPHTHALMOLOGY, (1986 Jan) 18 (1) 22-4. Journal code: 0210137. ISSN: 0003-4886.

L24 ANSWER 26 OF 31 MEDLINE

UPLICAT

TI Role of aldose reductase in the development of diabetes-associated complications.

SO AMERICAN JOURNAL OF MEDICINE, (1985 Nov 15) 79 (5A) 8-12

Journal code: 0267200. ISSN: 0002-9343.

L24 ANSWER 27 OF 31 MEDLINE

TI Morphologic effect of hyperosmolarity on rabbit corneal epithelium.

SO OPHTHALMOLOGY, (1984 Oct) 91 (10) 1205-12. Journal code: 7802443. ISSN: 0161-6420.

L24 ANSWER 28 OF 31 CAPLUS COPYRIGHT 2002 ACS

TI A model of epithelial water transport. The corneal endothelium

SO Biophys. J. (1981), 35(2), 315-38 CODEN: BIOJAU; ISSN: 0006-3495

L24 ANSWER 29 OF 31 CAPLUS COPYRIGHT 2002 ACS

TI Corneal pigment deposits from topically administered epinephrine. Experimental production

SO Arch. Ophthalmol. (1969), 82(6), 836-9 CODEN: AROPAW

L24 ANSWER 30 OF 31 MEDLINE

TI Anatomic study of water movement through rabbit corneal epithelium.

SO AMERICAN JOURNAL OF OPHTHALMOLOGY, (1969 Jan) 67 (1) 110-6.

Journal code: 0370500. ISSN: 0002-9394.

L24 ANSWER 31 OF 31 MEDLINE

TI The permeability of the corneal epithelium and endothelium to water.

SO EXPERIMENTAL EYE RESEARCH, (1967 Jan) 6 (1) 10-32. Journal code: 0370707. ISSN: 0014-4835.

=> D IBIB AB 27,25,22,21,16,15,2

L24 ANSWER 27 OF 31 MEDLINE

ACCESSION NUMBER: 85087229 MEDLINE

DOCUMENT NUMBER: 85087229 PubMed ID: 6514286
TITLE: Morphologic effect of hyperosmolarity on rabbit
corneal epithelium.

AUTHOR: Gilbard J P; Carter J B; Sang D N; Refojo M F;

Hanninen L

A; Kenyon K R

CONTRACT NUMBER: EY00327 (NEI)

EY03373 (NEI)

SOURCE: OPHTHALMOLOGY, (1984 Oct) 91 (10) 1205-12. Journal code: 7802443. ISSN: 0161-6420.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals ENTRY MONTH: 198502

ENTRY DATE: Entered STN: 19900320 Last Updated on STN: 19970203

Entered Medline: 19850219

AB The morphologic effect of hyperosmolarity, equivalent to that seen in the tear film of patients with keratoconjunctivitis sicca (KCS), on rabbit corneal epithelium in vitro and in vivo was studied. In the in vitro studies, corneal epithelium

was grown in explant cultures. Control tissue was cultured in a 307 mOsm/L

medium. Epithelium cultured in the 333, 361 and 363 mOsm/L media showed

decreased intercellular connections, blunting and loss of microplicae, disruptions in cell membranes and cellular swelling with decreased cytoplasmic density. In in vivo studies, corneas bathed in balanced

solutions (BSS) concentrated to 330, 360, or 407 mOsm/L showed increased

cell desquamation, and the cell changes observed at similar osmolarities

in the in vitro studies. The tear film osmolarities observed in KCS are sufficient to cause the corneal epithelial changes seen in patients with this disease.

L24 ANSWER 25 OF 31 MEDLINE DUPLICATE 6

ACCESSION NUMBER: 86157304 MEDLINE

DOCUMENT NUMBER: 86157304 PubMed ID: 3954296 TITLE: A comparison of therapeutic bandage lenses, tarsorrhaphy,

and antibiotic and hypertonic saline on corneal epithelial wound healing.

AUTHOR: Ali Z; Insler M S

SOURCE: ANNALS OF OPHTHALMOLOGY, (1986 Jan) 18

(1) 22-4.

Journal code: 0210137. ISSN: 0003-4886.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English
FILE SEGMENT: Priority Journals

ENTRY MONTH: 198604 ENTRY DATE: Entered STN: 19900321

Last Updated on STN: 19900321 Entered Medline: 19860403

AB Bandage lenses, a complete tarsorrhaphy, chloramphenicol 0.5%, and sodium

chloride 2% were tested for their effect on corneal
epithelial wound healing in rabbits. The healing rate was
decreased with bandage lenses and increased by tarsorrhaphy. There

significant effect on healing rate with either chloramphenicol or hypertonic saline.

L24 ANSWER 22 OF 31 CAPLUS COPYRIGHT 2002 ACS ACCESSION NUMBER: 1988:448412 CAPLUS

DOCUMENT NUMBER: 109:48412

TITLE: The effect of different solutions of glycerin on

corneal thickness and cell morphology studied by specular microscopy and scanning electron microscopy

AUTHOR(S): Cerulli, Luciano; Corsi, A.; Pocobelli, A.; Ricci, F.;

Caggiati, A.

CORPORATE SOURCE: Dep. Exp. Med., Univ. "G. D'Annunzio",

Rome, Italy

SOURCE: Ital. J. Ophthalmol. (1987), 1(2), 67-77

CODEN: ITJOE8
DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effect of the topical administration of different hyperosmolar solns. of glycerin on rabbit's eyes was detd.; a substantial redn. in corneal thickness with was obsd. with 15% and 30% glycerin. A biphasic

effect on corneal thickness was recorded using more concd. glycerin eyedrops. In these cases, a short period of corneal thinning was followed

by a longer phase of corneal swelling. Recovery to basic values

occurred

within 3-4 h. Specular microscopy and SEM revealed that high conens. of

glycerin induced "blebs" on the cell surface and cell membrane ruptures.

Measurements of corneal epithelial thickness suggested that the topical administration of the weaker solns. is ineffective in dehydrating corneal stroma; moreover the proven cytotoxicity of the stronger solns. contraindicates their clin. use.

L24 ANSWER 21 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1987:320163 BIOSIS DOCUMENT NUMBER: BA84:39670

TITLE: TOPICAL HYPEROSMOLAR SOLUTIONS IN THE REDUCTION

OF CORNEAL EDEMA.

AUTHOR(S): INSLER M S; BENEFIELD D W; ROSS E V CORPORATE SOURCE: LSU EYE CENT., 2020 GRAVIER ST., NEW ORLEANS, LA. 70112.

SOURCE: CLAO (CONTACT LENS ASSOC OPHTHALMOL)
J, (1987) 13 (3),

149-151.

CODEN: CLAJEU. ISSN: 0733-8902.

FILE SEGMENT: BA; OLD LANGUAGE: English

AB When the corneal epithelium and endothelium are stressed to their physiologic limits, due to mechanical or other stressors, the cornea swells from its normal state of 78% hydration.

assess the effect hypertonic agents might have in moving water out of the cornea and thereby decreasing corneal thickness, we compared

the effects of 2% and 5% (680 and 1720 mosm/L) sodium chloride drops on

rabbit eyes. Treatment with two drops of 2% NaCl solution resulted in a

decrease in corneal thickness of 18.5% over 90 minutes, while with the 5%

NaCl solution the cornea thinned by 29.9%. Both solutions appear beneficial in treating corneal edema, though neither dehydrating gent

produced a statistically significant reduction in corneal thickness (P = 0.07).

L24 ANSWER 16 OF 31 MEDLINE

ACCESSION NUMBER: 96285806 MEDLINE

DOCUMENT NUMBER: 96285806 PubMed ID: 8713925

TITLE: The effect of osmolality on the shedding rate of the corneal epithelium.

AUTHOR: Wilson G

CORPORATE SOURCE: Vision Science Research Center, School of Optometry,

University of Alabama at Birmingham 35294-4390, USA.

CONTRACT NUMBER: EY03039 (NEI)

SOURCE: CORNEA, (1996 May) 15 (3) 240-4.

Journal code: 8216186. ISSN: 0277-3740.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals ENTRY MONTH: 199611

ENTRY DATE: Entered STN: 19961219

Last Updated on STN: 19961219 Entered Medline: 19961107

AB It has been stated that in dry eye hyperosmolality damages the epithelium of the ocular surface by increasing the rate at which cells are

shed. To test this hypothesis, paired excised rabbit corneas were superfused with balanced salt solutions in which all conditions were seld

constant except the osmolality in contact with the epithelium. For a period of 400 min the epithelium of one of the corneas was exposed to one

of seven test osmolalities (200-425 mOsm/kg), whereas the other cornea was

used as a control (305 mOsm/kg). The number of cells shed from each

corneal surface was counted, and thickness changes in the epithelium and

stroma were measured. Only hypoosmotic solutions of 260 mOsm/kg or less

increased the number of cells shed relative to the control. None of the hyperosmotic solutions significantly increased the shedding rate. There was no significant change in the thickness of the epithelium, whereas the stroma swelled in hypoosmotic solutions and thinned in hyperosmotic solutions. It is concluded that hyperosmolality in the range encountered in dry eyes is not

sufficient in itself to increase the shedding rate.

L24 ANSWER 15 OF 31 MEDLINE

ACCESSION NUMBER: 97343625 MEDLINE

DOCUMENT NUMBER: 97343625 PubMed ID: 9200164

Sequential staining: the effects of sodium fluorescein, TITLE:

osmolarity, and pH on human corneal

epithelium.

Thomas M L; Szeto V R; Gan C M; Polse K A AUTHOR: CORPORATE SOURCE: University of California at Berkeley, School of Optometry,

USA.

CONTRACT NUMBER: EY-R01-07728 (NEI)

OPTOMETRY AND VISION SCIENCE, (1997 Apr) SOURCE:

74 (4) 207-10.

Journal code: 8904931. ISSN: 1040-5488.

United States PUB. COUNTRY:

Journal; Article; (JOURNAL ARTICLE) DOCUMENT TYPE:

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 199708

Entered STN: 19970825 ENTRY DATE: Last Updated on STN: 19980206 Entered Medline: 19970814

AB BACKGROUND: Previous reports have suggested that sequential applications

of sodium fluorescein (NaF) to the ocular surface cause loss of epithelial

cells. In those experiments the solutions were free of preservatives, but

delivered a hypertonic and alkaline load. It is possible that either the hyperosmolarity, NaF, and/or alkalinity may have contributed to the epithelial cell loss. Our study explored the possible impact of these three factors on epithelial integrity. METHODS: We

paradigm in which we designed four test solutions to isolate the cytotoxic

effects of the three factors. Fifteen subjects were exposed to one of the

solutions on separate visits. One solution was instilled in both eyes, every 3 min, for a total of seven applications, and a slitlamp examination

was performed. Staining was graded on a scale of 0 to 3 for each of

corneal sectors. RESULTS: Application of solution 1 (NaF, hyperosmotic, and alkaline) resulted in staining for all subjects. For the other three solutions (without NaF), insignificant staining occurred. CONCLUSIONS: The epithelium tolerates changes in pH and

osmolarity between 7.2 to 7.8 and 290 to 350, respectively. However,

when applied in multiple doses may be cytotoxic.

L24 ANSWER 2 OF 31 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2001:351674 BIOSIS DOCUMENT NUMBER: PREV200100351674

TITLE: Osmosensitive taurine transporter expression and uptake by

human corneal epithelial cells.

Shioda, R. (1); Reinach, P. S.; Kaminogawa, S.; AUTHOR(S):

Shimizu.

M.; Hisatsune, T. (1); Miyamoto, Y. (1)

CORPORATE SOURCE: (1) Integrated Biosciences, Univ of Tokyo, Bunkyo, Tokyo

Japan

IOVS, (March 15, 2001) Vol. 42, No. 4, pp. S588. SOURCE:

print.

Meeting Info.: Annual Meeting of the Association for Research in Vision and Ophthalmology Fort Lauderdale, Florida, USA April 29-May 04, 2001

DOCUMENT TYPE: Conference LANGUAGE: **English** SUMMARY LANGUAGE: English

=> d his

(FILE 'HOME' ENTERED AT 10:27:45 ON 01 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:28:13 ON 01 SEP 2002

4177 S DRY EYE OR XEROSIS OR HYPOLACRIMIA OR KERATOCONJUCTIVITIS SIC

251329 S ANIMAL MODEL L2

49 S L1 AND L2 L3

43 DUP REM L3 (6 DUPLICATES REMOVED) L4

L5 11300 S CORNEA? EPITHELI?

L6 631428 S DAMAGE

L7 708684 S DAMAG?

565797 S DESSICAT? OR DRY L8

L9 486 S L5(S)L6

L10 142 S L5(S)L8

L11 5 S L2 AND L10

5 DUP REM L11 (0 DUPLICATES REMOVED) L12

L13 586 S L5(S)L7

32 S L13 AND L8 L14

21 DUP REM L14 (11 DUPLICATES REMOVED)

130776 S OSMOT? OR OSMOS? L16

L17 14 S L9 AND L16

9 DUP REM L17 (5 DUPLICATES REMOVED) L18

29959 S HYPERTONIC L19

L20 8 S L1 AND L19

6 DUP REM L20 (2 DUPLICATES REMOVED) L21

L22 40993 S HYPERTONIC OR HYPEROSMO?

40 S L22 AND L5 L23

31 DUP REM L23 (9 DUPLICATES REMOVED) L24

=> log hold

L15

COST IN U.S. DOLLARS SINCE FILE TOTAL

SESSION ENTRY

FULL ESTIMATED COST 215.90 216.11

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE TOTAL

ENTRY SESSION

CA SUBSCRIBER PRICE

-1.24 -1.24

SESSION WILL BE HELD FOR 60 MINUTES STN INTERNATIONAL SESSION SUSPENDED AT 11:23:48 ON 01 SEP 2002

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1636DXS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * * * * * * Welcome to STN International * * * * * * * * *

Web Page URLs for STN Seminar Schedule - N. NEWS 1 America

NEWS 2 Apr 08 "Ask CAS" for self-help around the clock

NEWS 3 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area

NEWS 4 Apr 09 ZDB will be removed from STN

NEWS 5 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB

NEWS 6 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS

NEWS 7 Apr 22 BIOSIS Gene Names now available in TOXCENTER

NEWS 8 Apr 22 Federal Research in Progress (FEDRIP) now available

NEWS 9 Jun 03 New e-mail delivery for search results now available

NEWS 10 Jun 10 MEDLINE Reload

NEWS 11 Jun 10 PCTFULL has been reloaded

NEWS 12 Jul 02 FOREGE no longer contains STANDARDS file segment

NEWS 13 Jul 22 USAN to be reloaded July 28, 2002; saved answer sets no longer valid

NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY

NEWS 15 Jul 30 NETFIRST to be removed from STN

NEWS 16 Aug 08 CANCERLIT reload

NEWS 17 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN

NEWS 18 Aug 08 NTIS has been reloaded and enhanced

NEWS 19 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN

NEWS 20 Aug 19 IFIPAT, IFICDB, and IFIUDB have been reloaded NEWS 21 Aug 19 The MEDLINE file segment of TOXCENTER has been reloaded

NEWS 22 Aug 26 Sequence searching in REGISTRY enhanced NEWS 23 Sep 03 JAPIO has been reloaded and enhanced

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,

CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),

AND CURRENT DISCOVER FILE IS DATED 05

FEBRUARY 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS INTER General Internet Information

Welcome Banner and News Items NEWS LOGIN

NEWS PHONE Direct Dial and Telecommunication Network Access to STN

CAS World Wide Web Site (general information) **NEWS WWW**

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 10:20:10 ON 09 SEP 2002

=> file medline

SINCE FILE TOTAL COST IN U.S. DOLLARS ENTRY SESSION

FULL ESTIMATED COST 0.21 0.21

FILE 'MEDLINE' ENTERED AT 10:20:58 ON 09 SEP 2002

FILE LAST UPDATED: 7 SEP 2002 (20020907/UP). FILE COVERS

1958 TO DATE.

On June 9, 2002, MEDLINE was reloaded. See HELP RLOAD for details.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2002 vocabulary. Enter HELP THESAURUS for details.

THIS FILE CONTAINS CAS REGISTRY NUMBERS FOR EASY AND ACCURATE SUBSTANCE IDENTIFICATION.

=> file caplus

SINCE FILE TOTAL COST IN U.S. DOLLARS

ENTRY SESSION

FULL ESTIMATED COST

0.38 0.59

FILE 'CAPLUS' ENTERED AT 10:21:04 ON 09 SEP 2002 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 9 Sep 2002 VOL 137 ISS 11 FILE LAST UPDATED: 8 Sep 2002 (20020908/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or

the CAS Roles thesaurus (/RL field) in this file.

=> s ophthalmology and 91 and 1205/so

409 OPHTHALMOLOGY

1159 OPHTHALMOL

1421 OPHTHALMOLOGY

(OPHTHALMOLOGY OR OPHTHALMOL)

134822 91

4101 1205/SO

0 OPHTHALMOLOGY AND 91 AND 1205/SO L1

=> s ophthalmology/so and 91/so

1197 OPHTHALMOLOGY/SO

200651 91/SO

0 OPHTHALMOLOGY/SO AND 91/SO L2

=> s (morphologic effect of hyperosmolarity on rabbit corneal epithelium)/ti

1640 MORPHOLOGIC/TI

1 MORPHOLOGICS/TI

1641 MORPHOLOGIC/TI

((MORPHOLOGIC OR MORPHOLOGICS)/TI)

13 MORPHOL/TI

1654 MORPHOLOGIC/TI

((MORPHOLOGIC OR MORPHOL)/TI)

961668 EFFECT/TI

571823 EFFECTS/TI

1529188 EFFECT/TI

((EFFECT OR EFFECTS)/TI)

110 HYPEROSMOLARITY/TI

51533 RABBIT/TI
27607 RABBITS/TI
78938 RABBIT/TI
((RABBIT OR RABBITS)/TI)
3325 CORNEAL/TI
9478 EPITHELIUM/TI
8 EPITHELIUMS/TI
1672 EPITHELIA/TI
11136 EPITHELIUM/TI

((EPITHELIUM OR EPITHELIUMS OR EPITHELIA)/TI)
L3 0 (MORPHOLOGIC EFFECT OF HYPEROSMOLARITY
ON RABBIT CORNEAL EPITHEL

IUM)/TI

((MORPHOLOGIC(W)EFFECT(1W)HYPEROSMOLARITY(1W)RA BBIT(W)CORNEAL(W)EPITHELIUM/TI)

=> file medline biosis caplus

COST IN U.S. DOLLARS

SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST

19.49 20.08

FILE 'MEDLINE' ENTERED AT 10:23:27 ON 09 SEP 2002

FILE 'BIOSIS' ENTERED AT 10:23:27 ON 09 SEP 2002 COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'CAPLUS' ENTERED AT 10:23:27 ON 09 SEP 2002 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS. COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s ophthalmology/so and 1205/so

L4 34 OPHTHALMOLOGY/SO AND 1205/SO

=> s 14 and 91/so

L5 1 L4 AND 91/SO

=> d ti so 1

L5 ANSWER 1 OF 1 MEDLINE

TI Morphologic effect of hyperosmolarity on rabbit corneal epithelium.

SO OPHTHALMOLOGY, (1984 Oct) 91 (10) 1205-12.

Journal code: 7802443. ISSN: 0161-6420.

=> d ibib ab

L5 ANSWER 1 OF 1 MEDLINE

ACCESSION NUMBER: 85087229 MEDLINE

DOCUMENT NUMBER: 85087229 PubMed ID: 6514286

TITLE: Morphologic effect of hyperosmolarity on rabbit corneal epithelium.

AUTHOR: Gilbard J P; Carter J B; Sang D N; Refojo M F;

Hanninen L

A; Kenyon K R

CONTRACT NUMBER: EY00327 (NEI)

EY03373 (NEI)

SOURCE: OPHTHALMOLOGY, (1984 Oct) 91 (10)

1205-12.

Journal code: 7802443. ISSN: 0161-6420.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 198502

ENTRY DATE: Entered STN: 19900320

Last Updated on STN: 19970203 Entered Medline: 19850219

AB The morphologic effect of hyperosmolarity, equivalent to that seen

n the

tear film of patients with keratoconjunctivitis sicca (KCS), on rabbit

corneal epithelium in vitro and in vivo was studied. In the in vitro studies, corneal epithelium was grown in explant cultures. Control

was cultured in a 307 mOsm/L medium. Epithelium cultured in the 333, 361

and 363 mOsm/L media showed decreased intercellular connections, blunting

and loss of microplicae, disruptions in cell membranes and cellular swelling with decreased cytoplasmic density. In in vivo studies,

bathed in balanced salt solutions (BSS) concentrated to 330, 360, or

mOsm/L showed increased cell desquamation, and the cell changes observed

at similar osmolarities in the in vitro studies. The tear film osmolarities observed in KCS are sufficient to cause the corneal epithelial changes seen in patients with this disease.

=> s 15 <cit>

SmartSELECT INITIATED

New TRANSFER and ANALYZE Commands Now Available See HELP TRANSFER and HELP ANALYZE for Details

SET SMARTSELECT ON SET COMMAND COMPLETED

SEL L5 1- CIT

L6 SEL L5 1- CIT: 1 TERM

SET SMARTSELECT OFF SET COMMAND COMPLETED

S L6

'RE' IS NOT A VALID FIELD CODE 'RE' IS NOT A VALID FIELD CODE

7 1 L6

=> d ti so

L7 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2002 ACS

TI Comparison of hypotonic and isotonic solutions containing sodium hyaluronate on the symptomatic treatment of dry eye patients

SO Ophthalmologica (2001), 215(2), 124-127 CODEN: OPHTAD; ISSN: 0030-3755

=> d his

L1

(FILE 'HOME' ENTERED AT 10:20:10 ON 09 SEP 2002)

FILE 'MEDLINE' ENTERED AT 10:20:58 ON 09 SEP 2002

FILE 'CAPLUS' ENTERED AT 10:21:04 ON 09 SEP 2002

0 S OPHTHALMOLOGY AND 91 AND 1205/SO

L2 0 S OPHTHALMOLOGY/SO AND 91/SO

L3 0 S (MORPHOLOGIC EFFECT OF HYPEROSMOLARITY ON RABBIT CORNEAL EPIT

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 10:23:27 ON 09 SEP 2002

L4 34 S OPHTHALMOLOGY/SO AND 1205/SO

L5 1 S L4 AND 91/SO

SET SMARTSELECT ON

L6 SEL L5 1- CIT: 1 TERM

SET SMARTSELECT OFF

L7 1 S L6

=> log hold

COST IN U.S. DOLLARS SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST 18.90 38.98

SESSION WILL BE HELD FOR 60 MINUTES STN INTERNATIONAL SESSION SUSPENDED AT 10:25:39 ON 09 SEP 2002

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1636DXS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

******* Welcome to STN International *******

NEWS 1 Web Page URLs for STN Seminar Schedule - N.

America

NEWS 2 Apr 08 "Ask CAS" for self-help around the clock

NEWS 3 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area

NEWS 4 Apr 09 ZDB will be removed from STN

NEWS 5 Apr 19 US Patent Applications available in IFICDB,

IFIPAT, and IFIUDB

NEWS 6 Apr 22 Records from IP.com available in CAPLUS,

HCAPLUS, and ZCAPLUS

NEWS 7 Apr 22 BIOSIS Gene Names now available in

TOXCENTER

NEWS 8 Apr 22 Federal Research in Progress (FEDRIP) now available

NEWS 9 Jun 03 New e-mail delivery for search results now available

NEWS 10 Jun 10 MEDLINE Reload

NEWS 11 Jun 10 PCTFULL has been reloaded

NEWS 12 Jul 02 FOREGE no longer contains STANDARDS file segment

NEWS 13 Jul 22 USAN to be reloaded July 28, 2002; saved answer sets no longer valid

NEWS 14 Jul 29 Enhanced polymer searching in REGISTRY

NEWS 15 Jul 30 NETFIRST to be removed from STN

NEWS 16 Aug 08 CANCERLIT reload

NEWS 17 $\overline{\text{Aug}}$ 08 PHARMAMarketLetter(PHARMAML) - new on STN

NEWS 18 Aug 08 NTIS has been reloaded and enhanced

NEWS 19 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN

NEWS 20 Aug 19 IFIPAT, IFICDB, and IFIUDB have been reloaded

NEWS 21 $\overline{\text{Aug}}$ 19 The MEDLINE file segment of TOXCENTER has been reloaded

NEWS 22 Aug 26 Sequence searching in REGISTRY enhanced

NEWS 23 Sep 03 JAPIO has been reloaded and enhanced

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,

CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),

AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS INTER General Internet Information

NEWS LOGIN Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access

to STN

NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 12:07:17 ON 09 SEP 2002

=> file medline biosis caplus

COST IN U.S. DOLLARS

SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST

0.21 0.21

FILE 'MEDLINE' ENTERED AT 12:07:28 ON 09 SEP 2002

FILE 'BIOSIS' ENTERED AT 12:07:28 ON 09 SEP 2002 COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'CAPLUS' ENTERED AT 12:07:28 ON 09 SEP 2002 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP US AGETERMS" FOR DETAILS. COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s dry eye or keratoconjunctivitis sicca

L1 4126 DRY EYE OR KERATOCONJUNCTIVITIS SICCA

=> s screen

L2 125661 SCREEN

=> s 11 and 12

L3 6 L1 AND L2

=> dup rem 13

PROCESSING COMPLETED FOR L3

L4 3 DUP REM L3 (3 DUPLICATES REMOVED)

=> d ti so 1-3

L4 ANSWER 1 OF 3 MEDLINE

DUPLICATE 1

TI Air pollutants and tear film stability--a method for experimental evaluation.

SO CLINICAL PHYSIOLOGY, (2001 May) 21 (3) 282-6. Journal code: 8309768. ISSN: 0144-5979.

L4 ANSWER 2 OF 3 MEDLINE

TI Sensitivity and specificity of a screening questionnaire for dry eve.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1998) 438 807-20.

Journal code: 0121103. ISSN: 0065-2598.

L4 ANSWER 3 OF 3 MEDLINE

DUPLICATE 2

TI Assessment of a patient-completed questionnaire for keratoconjunctivitis sicca.

SO MEDICAL JOURNAL OF AUSTRALIA, (1988 Apr 18) 148 (8)

Journal code: 0400714. ISSN: 0025-729X.

=> s model

L5 3108267 MODEL

=> s 11 and 15

L6 220 L1 AND L5

=> s treat?

L7 6110040 TREAT?

=> s 16 and 17

L8 79 L6 AND L7

=> dup rem 18

PROCESSING COMPLETED FOR L8

L9

=> 19 and not py>2000

L9 IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system. For a list of commands available to you in the current file, enter "HELP COMMANDS" at an arrow prompt (=>).

=> s 19 and not py>2000 MISSING TERM 'AND NOT'

The search profile that was entered contains a logical operator followed immediately by another operator.

=> s l9 not py>2000 L10 35 L9 NOT PY>2000

=> d ti so 1-35

L10 ANSWER 1 OF 35 MEDLINE

TI [Use of keratoprosthesis type "champagne cork"--case report].

Zastosowanie keratoprotezy typu "korek od szampana"--opis
przypadku

SO KLINIKA OCZNA, (2000) 102 (5) 361-5. Journal code: 0376614. ISSN: 0023-2157.

L10 ANSWER 2 OF 35 MEDLINE

TI Microvascular submandibular gland transfer for severe cases of keratoconjunctivitis sicca.

SO PLASTIC AND RECONSTRUCTIVE SURGERY, (2000 Sep) 106

(3) 554-60; discussion

561-2.

Journal code: 1306050. ISSN: 0032-1052.

L10 ANSWER 3 OF 35 MEDLINE

TI Cyclosporine has a direct effect on the differentiation of a mucin-secreting cell line.

SO JOURNAL OF CELLULAR PHYSIOLOGY, (2000 Sep) 184 (3) 400-8.

Journal code: 0050222. ISSN: 0021-9541.

L10 ANSWER 4 OF 35 MEDLINE

TI Does androgen insufficiency cause lacrimal gland inflammation and aqueous

tear deficiency?.

SO INVESTIGATIVE OPHTHALMOLOGY AND VISUAL SCIENCE, (1999 May) 40 (6) 1261-5.

Journal code: 7703701. ISSN: 0146-0404.

L10 ANSWER 5 OF 35 MEDLINE

TI Ocular contact time of a carbomer gel (GelTears) in humans.

SO BRITISH JOURNAL OF OPHTHALMOLOGY, (1998 Oct) 82 (10) 1131-4.

Journal code: 0421041. ISSN: 0007-1161.

L10 ANSWER 6 OF 35 MEDLINE

TI [Important results of about 45 years balneo-medical research in Bad Hall].

Wichtige Ergebnisse aus rund 45 Jahren balneomedizinischer Forschung in

Bad Hall.

SO WIENER MEDIZINISCHE WOCHENSCHRIFT. SUPPLEMENT, (1998) 148 (110) 3-11.

Ref: 70

Journal code: 0413007. ISSN: 0301-7826.

L10 ANSWER 7 OF 35 MEDLINE

TI [Keratoconjunctivitis sicca in Sjogren's syndrome: diagnostic significance of relative protein composition of tears]. Sukhoi keratokon"iunktivit pri sindrome Shegrena: diagnosticheskoe znachenie izmeneniia otnositel'nogo belkovogo sostava slezy.

SO VESTNIK OFTALMOLOGII, (1998 Mar-Apr) 114 (2) 40-2. Journal code: 0415216. ISSN: 0042-465X.

L10 ANSWER 8 OF 35 MEDLINE

TI Lactoferrin suppresses loss of corneal epithelial integrity in a rabbit short-term dry eye model.

SO JOURNAL OF OCULAR PHARMACOLOGY AND

THERAPEUTICS, (1998 Apr.) 14 (2) 99-107.

Journal code: 9511091. ISSN: 1080-7683.

L10 ANSWER 9 OF 35 MEDLINE

TI Androgen stimulation of lacrimal gland function in mouse models of Sjogren's syndrome.

SO JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY, (1997 Feb) 60 (3-4)

237-45.

Journal code: 9015483. ISSN: 0960-0760.

L10 ANSWER 10 OF 35 MEDLINE

TI Clinical features, pathogenesis, and treatment of Sjogren's syndrome.

SO CURRENT OPINION IN RHEUMATOLOGY, (1996 Sep) 8 (5) 438-45. Ref: 104

Journal code: 9000851. ISSN: 1040-8711.

L10 ANSWER 11 OF 35 MEDLINE

TI Establishment of a rabbit short-term dry eye model.

SO JOURNAL OF OCULAR PHARMACOLOGY AND

THERAPEUTICS, (1995 Winter) 11 (4)

Journal code: 9511091. ISSN: 1080-7683.

L10 ANSWER 12 OF 35 MEDLINE

TI Dry eye with only decreased tear break-up time is sometimes associated with allergic conjunctivitis.

SO OPHTHALMOLOGY, (1995 Feb) 102 (2) 302-9. Journal code: 7802443. ISSN: 0161-6420.

L10 ANSWER 13 OF 35 MEDLINE

TI Epidemiology, pathogenesis, animal models, and treatment of Sjogren's syndrome.

SO CURRENT OPINION IN RHEUMATOLOGY, (1994 Sep) 6 (5)

501-8. Ref: 89

Journal code: 9000851. ISSN: 1040-8711.

L10 ANSWER 14 OF 35 MEDLINE

TI Treatment of keratoconjunctivitis sicca in

rabbits with 3-isobutyl-1-methylxanthine.

SO ARCHIVES OF OPHTHALMOLOGY, (1994 Dec) 112 (12) 1614-6.

Journal code: 7706534. ISSN: 0003-9950.

L10 ANSWER 15 OF 35 MEDLINE

TI Characteristics of a canine model of KCS: effective treatment with topical cyclosporine.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1994) 350 583-94. Ref: 56

Journal code: 0121103. ISSN: 0065-2598.

L10 ANSWER 16 OF 35 MEDLINE

TI Collagen-based drug delivery and artificial tears.

SO JOURNAL OF OCULAR PHARMACOLOGY, (1994 Spring) 10 (1) 17-27.

Journal code: 8511297. ISSN: 8756-3320.

L10 ANSWER 17 OF 35 MEDLINE

TI Characterization of water retentive properties of hyaluronan.

SO CORNEA, (1993 Sep) 12 (5) 433-6. Journal code: 8216186. ISSN: 0277-3740.

L10 ANSWER 18 OF 35 MEDLINE

TI Efficacy of laser punctal occlusion.

SO OPHTHALMOLOGY, (1992 Apr) 99 (4) 618-21. Journal code: 7802443. ISSN: 0161-6420.

L10 ANSWER 19 OF 35 MEDLINE

TI An electrolyte-based solution that increases comeal glycogen and

conjunctival goblet-cell density in a rabbit model for keratoconjunctivitis sicca.

SO OPHTHALMOLOGY, (1992 Apr) 99 (4) 600-4. Journal code: 7802443. ISSN: 0161-6420.

L10 ANSWER 20 OF 35 MEDLINE

TI Beneficial effects of a retinoic acid analog, CBS-211 A, on an experimental model of keratoconjunctivitis sicca.

SO INVESTIGATIVE OPHTHALMOLOGY AND VISUAL SCIENCE, (1992 Jan) 33 (1) 190-5.

Journal code: 7703701. ISSN: 0146-0404.

L10 ANSWER 21 OF 35 MEDLINE

TI Eledoisin and lacrimal secretion in the rabbit.

SO CURRENT EYE RESEARCH, (1990 Mar) 9 (3) 273-6. Journal code: 8104312. ISSN: 0271-3683.

L10 ANSWER 22 OF 35 MEDLINE

TI Spontaneous canine keratoconjunctivitis sicca. A useful model for human keratoconjunctivitis sicca: treatment with cyclosporine eye drops.

SO ARCHIVES OF OPHTHALMOLOGY, (1989 Aug) 107 (8) 1210-

Journal code: 7706534. ISSN: 0003-9950.

L10 ANSWER 23 OF 35 MEDLINE

TI Keratoconjunctivitis sicca in the dog following 5-aminosalicylic acid administration.

SO HUMAN TOXICOLOGY, (1987 Sep) 6 (5) 377-83. Journal code: 8206759. ISSN: 0144-5952.

L10 ANSWER 24 OF 35 MEDLINE

TI Tear physiology and dry eyes.

SO SURVEY OF OPHTHALMOLOGY, (1977 Sep-Oct) 22 (2) 69-87. Ref: 119

Journal code: 0404551. ISSN: 0039-6257.

L10 ANSWER 25 OF 35 MEDLINE

TI Keratoconjunctivitis sicca.

SO TRANSACTIONS - AMERICAN ACADEMY OF OPHTHALMOLOGY AND OTOLARYNGOLOGY, (1976

Jul-Aug) 81 (4 pt 1) OP619-23.

Journal code: 7506085. ISSN: 0002-7154.

L10 ANSWER 26 OF 35 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Development of a simple dry eye model in the albino rabbit and evaluation of some tear substitutes.

SO Ophthalmic Research, (May-June, 1999) Vol. 31, No. 3, pp. 229-235.

ISSN: 0030-3747.

L10 ANSWER 27 OF 35 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The effects of artificial tear solutions on wound healing in full thickness comeal incisions.

SO Acta Physiologica Hungarica, (1997) Vol. 85, No. 3, pp. 251-258. ISSN: 0231-424X.

L10 ANSWER 28 OF 35 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI The natural history of dry-eye disease: Implications for diagnosis and treatment.

SO Journal of Rheumatology, (1997) Vol. 24, No. SUPPL. 50, pp. 49. Meeting Info.: VIth International Symposium on Sjogren's Syndrome

Connecticut, USA October 15-18, 1997

ISSN: 0315-162X.

L10 ANSWER 29 OF 35 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

TI Characteristics of a canine model of KCS: Effective treatment with topical cyclosporine.

SO Sullivan, D. A. [Editor]. Advances in Experimental Medicine and Biology,

(1994) Vol. 350, pp. 583-594. Advances in Experimental Medicine and

Biology; Lacrimal gland, tear film, and dry eye syndromes: Basic science

and clinical relevance.

Publisher: Plenum Press 233 Spring Street, New York, New York,

Meeting Info.: International Conference Southampton, Bermuda November

14-17, 1992

ISSN: 0065-2598. ISBN: 0-306-44676-6.

L10 ANSWER 30 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI Dry eye treatment with eye drops that stimulate mucin production

SO Advances in Experimental Medicine and Biology (1998), 438(Lacrimal Gland,

Tear Film, and Dry Eye Syndromes 2), 965-968 CODEN: AEMBAP; ISSN: 0065-2598

L10 ANSWER 31 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI Gender and androgen treatment influence the expression of proto-oncogenes and apoptotic factors in lacrimal and salivary tissues

MRL/lpr mice

SO Clinical Immunology and Immunopathology (1998), 86(1), 59-71 CODEN: CLIIAT; ISSN: 0090-1229

L10 ANSWER 32 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI An in vitro technique for measuring contact angles on the corneal surface

and its application to evaluate corneal wetting properties of water soluble polymers

SO International Journal of Pharmaceutics (1995), 119(2), 149-63 CODEN: IJPHDE; ISSN: 0378-5173

L10 ANSWER 33 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI Preparation of polypeptide-polymer ion conjugates for treatment of wounds

SO Jpn. Kokai Tokkyo Koho, 12 pp. CODEN: JKXXAF

L10 ANSWER 34 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI Dehydroepiandrosterone compositions and therapy for dry eye syndrome

SO PCT Int. Appl., 30 pp CODEN: PIXXD2

L10 ANSWER 35 OF 35 CAPLUS COPYRIGHT 2002 ACS

TI Ocular androgen therapy in Sjogren's syndrome

SO PCT Int. Appl., 33 pp. CODEN: PIXXD2

=> d ibib ab 30,26,22,20,19,14

L10 ANSWER 30 OF 35 CAPLUS COPYRIGHT 2002 ACS

1998:562115 CAPLUS ACCESSION NUMBER: 129:310846

DOCUMENT NUMBER: TITLE:

Dry eye treatment with

eye drops that stimulate mucin production

AUTHOR(S): Hamano, Takashi

CORPORATE SOURCE: Dep. Ophthalmol., Osaka Univ. Med. Sch., Suita, Osaka,

Japan

SOURCE:

Advances in Experimental Medicine and Biology

(1998),

438(Lacrimal Gland, Tear Film, and Dry Eye Syndromes 2), 965-968

CODEN: AEMBAP; ISSN: 0065-2598

PUBLISHER:

Plenum Publishing Corp.

DOCUMENT TYPE:

Journal

LANGUAGE: English

AB In conjunctival goblet cells, 1.0% gefarnate eye drops increased the d. of

cells. In the rabbit dry eye model

treated with the gefarnate eye drops, there was less staining, indicating that desiccation of the comea caused less damage. Thus, gefarnate-contg. eye drops stimulate the expression of

transmembrane mucin

and stimulate mucin prodn. enough to decrease the corneal epithelial damage of dry eye.

L10 ANSWER 26 OF 35 BIOSIS COPYRIGHT 2002 BIOLOGICAL

ABSTRACTS INC. ACCESSION NUMBER: 1999:307975 BIOSIS DOCUMENT NUMBER: PREV199900307975 Development of a simple dry eye TITLE:

model in the albino rabbit and evaluation of some tear substitutes.

Burgalassi, Susi (1); Panichi, Luana; Chetoni, AUTHOR(S): Patrizia;

Saettone, M. Fabrizio; Boldrini, Enrico

CORPORATE SOURCE: (1) Department of Pharmaceutical Sciences, University of

Pisa, Via Bonanno 33, I-56126, Pisa Italy

Ophthalmic Research, (May-June, 1999) Vol. 31, No. SOURCE:

3, pp.

229-235.

ISSN: 0030-3747. DOCUMENT TYPE: Article LANGUAGE: English SUMMARY LANGUAGE: English

AB The present paper is concerned with the development of a simple dry eye model in the rabbit, induced by daily

repeated instillations of 1.0% atropine sulphate. The evolution of the dry eye syndrome in the animals was assessed by the Schirmer I test and by examination of the cornea after fluorescein staining. The model produced rapidly some typical dry eye symptoms and could be satisfactorily used for a preliminary assessment of the protective activity of some polymeric tear substitutes.

These were based on hydroxypropylmethylcellulose, sodium hyaluronate,

sodium polyacrylate or tamarind gum. The latter polymer showed the

overall results. Ferning tests on the formulations were also

their validity as predictors of the efficacy of tear substitutes is discussed.

L10 ANSWER 22 OF 35 MEDLINE

ACCESSION NUMBER: 89334669 MEDLINE

DOCUMENT NUMBER: 89334669 PubMed ID: 2757551

Spontaneous canine keratoconjunctivitis TITLE:

sicca. A useful model for human keratoconjunctivitis sicca:

treatment with cyclosporine eye drops.

Kaswan R L; Salisbury M A; Ward D A AUTHOR: CORPORATE SOURCE: Department of Small Animal Medicine, College of Veterinary

Medicine, University of Georgia, Athens 30602.

ARCHIVES OF OPHTHALMOLOGY, (1989 Aug) SOURCE: 107 (8) 1210-6.

Journal code: 7706534. ISSN: 0003-9950.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

Abridged Index Medicus Journals; Priority FILE SEGMENT:

Journals

ENTRY MONTH: 198909

Entered STN: 19900309 ENTRY DATE: Last Updated on STN: 19900309 Entered Medline: 19890901

AB Thirty-six sequential cases of canine keratoconjunctivitis sicca (KCS) were treated with ophthalmic cyclosporine.

The effects of topical cyclosporine were twofold: (1) cyclosporine increased tear production by 5 mm/min or greater in all cases of spontaneous KCS having an initial Schirmer's Tear Test value greater than

2 mm/min and in 59% of eyes with an initial Schirmer's Tear Test value of

0 to 2 mm/min, and (2) cyclosporine caused marked regression of chronic

corneal neovascularization and granulation even in eyes in which lacrimation failed to improve. Additional benefits of topical cyclosporine

were reduced mucopurulent conjunctivitis, rapid healing of nonhealing

corneal ulcers, and reduced dependence on frequent topical treatments of KCS. Twelve normal beagles treated with topical cyclosporine also had a reversible increase in lacrimation compared with baseline or placebo control-treated dogs.

L10 ANSWER 20 OF 35 MEDLINE

ACCESSION NUMBER: 92112433 MEDLINE

DOCUMENT NUMBER: 92112433 PubMed ID: 1730541

Beneficial effects of a retinoic acid analog, CBS-211 A, TITLE: on

an experimental model of

keratoconjunctivitis sicca.

Driot JY; Bonne C AUTHOR:

CORPORATE SOURCE: Chauvin Laboratory Research Centre,

Montpellier, France.

INVESTIGATIVE OPHTHALMOLOGY AND SOURCE:

VISUAL SCIENCE, (1992 Jan) 33 (1) 190-5.

Journal code: 7703701. ISSN: 0146-0404.

United States PUB. COUNTRY:

Journal; Article; (JOURNAL ARTICLE) DOCUMENT TYPE:

LANGUAGE: English

Priority Journals FILE SEGMENT: 199202 ENTRY MONTH:

Entered STN: 19920308 ENTRY DATE:

Last Updated on STN: 19920308 Entered Medline: 19920220

AB We report the effects of CBS-211 A, a synthetic retinoic acid analog, on a

previously described experimental model of

keratoconjunctivitis sicca (KCS) in the rabbit. A 9-week topical treatment with 0.02% CBS-211 A in aqueous vehicle significantly increased the conjunctival goblet cell density (P less

0.01, impression cytology counting), stopped the evolution of the corneo-conjunctival surface alteration (P less than 0.05, rose bengal test), and restored a basically normal mucosecretory product quality

goblet cells (lectin histochemistry) compared to vehicle treatment . The results assess the efficacy of this compound in reversing KCS pathology in a relevant model different from general vitamin A deficiency models, and strongly support the rationale for using such a well-tolerated retinoid in dry eye treatment.

L10 ANSWER 19 OF 35 MEDLINE

ACCESSION NUMBER: 92262107 MEDLINE

DOCUMENT NUMBER: 92262107 PubMed ID: 1584579 An electrolyte-based solution that increases corneal TITLE:

glycogen and conjunctival goblet-cell density in a rabbit model for keratoconjunctivitis

sicca.

Gilbard JP; Rossi SR AUTHOR:

CORPORATE SOURCE: Cornea Research Unit, Eye Research Institute, Boston, MA

02114.

CONTRACT NUMBER: EY-03373 (NEI)

OPHTHALMOLOGY, (1992 Apr) 99 (4) 600-4. SOURCE:

Journal code: 7802443. ISSN: 0161-6420.

PUB. COUNTRY: United States

Journal; Article; (JOURNAL ARTICLE) DOCUMENT TYPE:

LANGUAGE: English FILE SEGMENT: **Priority Journals**

ENTRY MONTH: 199206 Entered STN: 19920626 ENTRY DATE:

Last Updated on STN: 19920626 Entered Medline: 19920615

AB Thirty-two rabbits with monocular surgically induced keratoconjunctivitis sicca (KCS) underwent masked treatment for 12 weeks with 1 of 4 artificial tear solutions. Disease in each group of treated rabbits was compared with disease in untreated KCS controls. One of the solutions tested was a unique electrolyte-based formulation shown previously to preserve normal

goblet-cell density after extended exposure in normal rabbits. Only the

electrolyte-based solution decreased elevated tear osmolarity and sodium

after 9 weeks of treatment (P less than 0.05). At 20 weeks, mean corneal glycogen and conjunctival goblet-cell density in eyes treated with the electrolyte-based solution increased significantly relative to untreated KCS controls (P less than 0.01).

the other three solutions, mean glycogen levels and goblet-cell densities

were either decreased relative to untreated KCS controls (P less than 0.05) or were unchanged. The electrolyte-based solution is the first treatment to increase corneal glycogen and conjunctival goblet cells in a rabbit model of KCS.

L10 ANSWER 14 OF 35 MEDLINE

ACCESSION NUMBER: 95085434 MEDLINE

DOCUMENT NUMBER: 95085434 PubMed ID: 7527631

TITLE: Treatment of keratoconjunctivitis

sicca in rabbits with 3-isobutyl-1-methylxanthine.

AUTHOR: Gilbard J P

CORPORATE SOURCE: Schepens Eye Research Institute,

Department of

Ophthalmology, Harvard Medical School, Boston, Mass.

CONTRACT NUMBER: EYO3373 (NEI)

ARCHIVES OF OPHTHALMOLOGY, (1994 Dec) SOURCE:

112 (12) 1614-6.

Journal code: 7706534. ISSN: 0003-9950.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Abridged Index Medicus Journals; Priority

Journals

ENTRY MONTH: 199501

ENTRY DATE: Entered STN: 19950124 Last Updated on STN: 19960129 Entered Medline: 19950110

AB OBJECTIVE: To examine the effects of topical 3-isobutyl-1methylxanthine

treatment on tear-film osmolarity, conjunctival goblet-cell densities, and corneal epithelial glycogen levels in a rabbit model for keratoconjunctivitis sicca. METHODS:

Keratoconjunctivitis sicca was surgically induced in the right eyes of 16 rabbits. In a masked protocol, eight of these operated-on

eyes underwent treatment for 12 weeks with a 3.0-mmol solution of 3-isobutyl-1-methylxanthine. The remaining eight operated-on

left untreated and served as controls. RESULTS: The 3-isobutyl-1methylxanthine treatment resulted in a rapid and significant decrease in tear osmolarity and sodium (P < .5) and potassium levels (P <

.05) and a significant increase in conjunctival goblet-cell densities and

corneal epithelial glycogen levels compared with untreated and operated-on

controls (P < .001). CONCLUSIONS: 3-Isobutyl-1-methylxanthine

significantly decreased tear-film osmolarity in this rabbit model for keratoconjunctivitis sicca and restored

conjunctival goblet-cell densities and corneal glycogen levels, thus reversing the disease process.

=> d his

(FILE 'HOME' ENTERED AT 12:07:17 ON 09 SEP 2002)

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 12:07:28 ON 09 SEP 2002

Ll 4126 S DRY EYE OR KERATOCONJUNCTIVITIS SICCA

L2 125661 S SCREEN

L3 6 S L1 AND L2

L4 3 DUP REM L3 (3 DUPLICATES REMOVED)

L5 3108267 S MODEL

L6 220 S L1 AND L5

L7 6110040 S TREAT?

L8 79 S L6 AND L7

1.9 50 DUP REM L8 (29 DUPLICATES REMOVED)

L10 35 S L9 NOT PY>2000

=> log hold

COST IN U.S. DOLLARS

SINCE FILE TOTAL **ENTRY** SESSION

FULL ESTIMATED COST

42.13 42.34

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE TOTAL

ENTRY SESSION

CA SUBSCRIBER PRICE

-0.62 -0.62

SESSION WILL BE HELD FOR 60 MINUTES STN INTERNATIONAL SESSION SUSPENDED AT 12:19:04 ON 09 SEP 2002